

Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using lepton + jets events with lifetime b -tagging

DØ Collaboration

V.M. Abazov ^{am}, B. Abbott ^{bz}, M. Abolins ^{bq}, B.S. Acharya ^{ag}, M. Adams ^{bd}, T. Adams ^{bb}, M. Agelou ^u, J.-L. Agram ^{v,w}, S.H. Ahn ^{ai}, M. Ahsan ^{bk}, G.D. Alexeev ^{am}, G. Alkhazov ^{aq}, A. Alton ^{bp}, G. Alverson ^{bo}, G.A. Alves ^b, M. Anastasioie ^{al}, T. Andeen ^{bf}, S. Anderson ^{ax}, B. Andrieu ^t, Y. Arnoud ^q, A. Askew ^{bb}, B. Åsman ^{ar,as,at}, A.C.S. Assis Jesus ^c, O. Atramentov ^{bi}, C. Autermann ^y, C. Avila ^k, F. Badaud ^p, A. Baden ^{bm}, B. Baldin ^{bc}, P.W. Balm ^{ak}, S. Banerjee ^{ag}, E. Barberis ^{bo}, P. Bargassa ^{cd}, P. Baringer ^{bj}, C. Barnes ^{av}, J. Barreto ^b, J.F. Bartlett ^{bc}, U. Bassler ^t, D. Bauer ^{bg}, A. Bean ^{bj}, S. Beauceron ^t, M. Begalli ^c, M. Begel ^{bv}, A. Bellavance ^{bs}, S.B. Beri ^{ae}, G. Bernardi ^t, R. Bernhard ^{bc,1}, I. Bertram ^{au}, M. Besançon ^u, R. Beuselinck ^{av}, V.A. Bezzubov ^{ap}, P.C. Bhat ^{bc}, V. Bhatnagar ^{ae}, M. Binder ^{ac}, C. Biscarat ^{au}, K.M. Black ^{bn}, I. Blackler ^{av}, G. Blazey ^{be}, F. Blekman ^{av}, S. Blessing ^{bb}, D. Bloch ^{v,w}, U. Blumenschein ^{aa}, A. Boehnlein ^{bc}, O. Boeriu ^{bh}, T.A. Bolton ^{bk}, F. Borcherding ^{bc}, G. Borissov ^{au}, K. Bos ^{ak}, T. Bose ^{bu}, A. Brandt ^{cb}, R. Brock ^{bq}, G. Brooijmans ^{bu}, A. Bross ^{bc}, N.J. Buchanan ^{bb}, D. Buchholz ^{bf}, M. Buehler ^{bd}, V. Buescher ^{aa}, S. Burdin ^{bc}, S. Burke ^{ax}, T.H. Burnett ^{cf}, E. Busato ^t, C.P. Buszello ^{av}, J.M. Butler ^{bn}, J. Cammin ^{bv}, S. Caron ^{ak}, W. Carvalho ^c, B.C.K. Casey ^{ca}, N.M. Cason ^{bh}, H. Castilla-Valdez ^{aj}, S. Chakrabarti ^{ag}, D. Chakraborty ^{be}, K.M. Chan ^{bv}, A. Chandra ^{ag}, D. Chapin ^{ca}, F. Charles ^{v,w}, E. Cheu ^{ax}, D.K. Cho ^{bn}, S. Choi ^{ba}, B. Choudhary ^{af}, T. Christiansen ^{ac}, L. Christofek ^{bj}, D. Claes ^{bs}, B. Clément ^{v,w}, C. Clément ^{ar,as,at}, Y. Coadou ^{e,f,g,h}, M. Cooke ^{cd}, W.E. Cooper ^{bc}, D. Coppage ^{bj}, M. Corcoran ^{cd}, A. Cothenet ^r, M.-C. Cousinou ^r, B. Cox ^{aw}, S. Crépé-Renaudin ^q, D. Cutts ^{ca}, H. da Motta ^b, M. Das ^{bl}, B. Davies ^{au}, G. Davies ^{av}, G.A. Davis ^{bf}, K. De ^{cb}, P. de Jong ^{ak}, S.J. de Jong ^{al}, E. De La Cruz-Burelo ^{bp}, C. De Oliveira Martins ^c, S. Dean ^{aw}, J.D. Degenhardt ^{bp}, F. Déliot ^u, M. Demarteau ^{bc}, R. Demina ^{bv}, P. Demine ^u, D. Denisov ^{bc}, S.P. Denisov ^{ap}, S. Desai ^{bw}, H.T. Diehl ^{bc}, M. Diesburg ^{bc}, M. Doidge ^{au}, H. Dong ^{bw}, S. Doulas ^{bo}, L.V. Dudko ^{ao}, L. Duflot ^s, S.R. Dugad ^{ag}, A. Duperrin ^r, J. Dyer ^{bq}, A. Dyshkant ^{be},

M. Eads ^{be}, D. Edmunds ^{bq}, T. Edwards ^{aw}, J. Ellison ^{ba}, J. Elmsheuser ^{ac}, V.D. Elvira ^{bc}, S. Eno ^{bm}, P. Ermolov ^{ao}, O.V. Eroshin ^{ap}, J. Estrada ^{bc}, H. Evans ^{bu}, A. Evdokimov ^{an}, V.N. Evdokimov ^{ap}, J. Fast ^{bc}, S.N. Fatachia ^{bn}, L. Feligioni ^{bn}, A.V. Ferapontov ^{ap}, T. Ferbel ^{bv}, F. Fiedler ^{ac}, F. Filthaut ^{al}, W. Fisher ^{bt}, H.E. Fisk ^{bc}, I. Fleck ^{aa}, M. Fortner ^{be}, H. Fox ^{aa}, S. Fu ^{bc}, S. Fuess ^{bc}, T. Gadfort ^{cf}, C.F. Galea ^{al}, E. Gallas ^{bc}, E. Galyaev ^{bh}, C. Garcia ^{bv}, A. Garcia-Bellido ^{cf}, J. Gardner ^{bj}, V. Gavrilov ^{an}, A. Gay ^{v,w}, P. Gay ^p, D. Gelé ^{v,w}, R. Gelhaus ^{ba}, K. Genser ^{bc}, C.E. Gerber ^{bd}, Y. Gershtein ^{bb}, D. Gillberg ^{e,f,g,h}, G. Ginther ^{bv}, T. Golling ^z, N. Gollub ^{ar,as,at}, B. Gómez ^k, K. Gounder ^{bc}, A. Goussiou ^{bh}, P.D. Grannis ^{bw}, S. Greder ^c, H. Greenlee ^{bc}, Z.D. Greenwood ^{bl}, E.M. Gregores ^d, Ph. Gris ^p, J.-F. Grivaz ^s, L. Groer ^{bu}, S. Grünendahl ^{bc}, M.W. Grünewald ^{ah}, S.N. Gurzhiev ^{ap}, G. Gutierrez ^{bc}, P. Gutierrez ^{bz}, A. Haas ^{bu}, N.J. Hadley ^{bm}, S. Hagopian ^{bb}, I. Hall ^{bz}, R.E. Hall ^{az}, C. Han ^{bp}, L. Han ^j, K. Hanagaki ^{bc}, K. Harder ^{bk}, A. Harel ^{ad}, R. Harrington ^{bo}, J.M. Hauptman ^{bi}, R. Hauser ^{bq}, J. Hays ^{bf}, T. Hebbeker ^y, D. Hedin ^{be}, J.M. Heinmiller ^{bd}, A.P. Heinson ^{ba}, U. Heintz ^{bn}, C. Hensel ^{bj}, G. Hesketh ^{bo}, M.D. Hildreth ^{bh}, R. Hirosky ^{ce}, J.D. Hobbs ^{bw}, B. Hoeneisen ^o, M. Hohlfeld ^{ab}, S.J. Hong ^{ai}, R. Hooper ^{ca}, P. Houben ^{ak}, Y. Hu ^{bw}, J. Huang ^{bg}, V. Hynek ^l, I. Iashvili ^{ba}, R. Illingworth ^{bc}, A.S. Ito ^{bc}, S. Jabeen ^{bj}, M. Jaffré ^s, S. Jain ^{bz}, V. Jain ^{bx}, K. Jakobs ^{aa}, A. Jenkins ^{av}, R. Jesik ^{av}, K. Johns ^{ax}, M. Johnson ^{bc}, A. Jonckheere ^{bc}, P. Jonsson ^{av}, A. Juste ^{bc}, M.M. Kado ^{ay}, D. Käfer ^y, S. Kahn ^{bx}, E. Kajfasz ^r, A.M. Kalinin ^{am}, J. Kalk ^{bq}, D. Karmanov ^{ao}, J. Kasper ^{bn}, D. Kau ^{bb}, R. Kaur ^{ae}, R. Kehoe ^{cc}, S. Kermiche ^r, S. Kesisoglou ^{ca}, A. Khanov ^{bv}, A. Kharchilava ^{bh}, Y.M. Kharzeev ^{am}, H. Kim ^{cb}, T.J. Kim ^{ai}, B. Klima ^{bc}, M. Klute ^z, J.M. Kohli ^{ae}, J.-P. Konrath ^{aa}, M. Kopal ^{bz}, V.M. Korablev ^{ap}, J. Kotcher ^{bx}, B. Kothari ^{bu}, A. Koubarovsky ^{ao}, A.V. Kozelov ^{ap}, J. Kozminski ^{bq}, A. Kryemadhi ^{ce}, S. Krzywdzinski ^{bc}, Y. Kulik ^{bc}, A. Kumar ^{af}, S. Kunori ^{bm}, A. Kupco ⁿ, T. Kurča ^x, J. Kvita ^l, S. Lager ^{ar,as,at}, N. Lahrichi ^u, G. Landsberg ^{ca}, J. Lazoflores ^{bb}, A.-C. Le Bihan ^{v,w}, P. Lebrun ^x, W.M. Lee ^{bb}, A. Leflat ^{ao}, F. Lehner ^{bc,l}, C. Leonidopoulos ^{bu}, J. Leveque ^{ax}, P. Lewis ^{av}, J. Li ^{cb}, Q.Z. Li ^{bc}, J.G.R. Lima ^{be}, D. Lincoln ^{bc}, S.L. Linn ^{bb}, J. Linnemann ^{bq}, V.V. Lipaev ^{ap}, R. Lipton ^{bc}, L. Lobo ^{av}, A. Lobodenko ^{aq}, M. Lokajicek ⁿ, A. Lounis ^{v,w}, P. Love ^{au}, H.J. Lubatti ^{cf}, L. Lueking ^{bc}, M. Lynker ^{bh}, A.L. Lyon ^{bc}, A.K.A. Maciel ^{be}, R.J. Madaras ^{ay}, P. Mättig ^{ad}, C. Magass ^y, A. Magerkurth ^{bp}, A.-M. Magnan ^q, N. Makovec ^s, P.K. Mal ^{ag}, H.B. Malbouisson ^c, S. Malik ^{bs}, V.L. Malyshev ^{am}, H.S. Mao ⁱ, Y. Maravin ^{bc}, M. Martens ^{bc}, S.E.K. Mattingly ^{ca}, A.A. Mayorov ^{ap}, R. McCarthy ^{bw}, R. McCroskey ^{ax}, D. Meder ^{ab}, A. Melnitchouk ^{br}, A. Mendes ^r, M. Merkin ^{ao}, K.W. Merritt ^{bc}, A. Meyer ^y, J. Meyer ^z, M. Michaut ^u, H. Miettinen ^{cd}, J. Mitrevski ^{bu}, J. Molina ^c, N.K. Mondal ^{ag}, R.W. Moore ^{e,f,g,h}, T. Moulik ^{bj}, G.S. Muanza ^x, M. Mulders ^{bc}, L. Mundim ^c, Y.D. Mutaf ^{bw}, E. Nagy ^r, M. Narain ^{bn}, N.A. Naumann ^{al}, H.A. Neal ^{bp}, J.P. Negret ^k, S. Nelson ^{bb}, P. Neustroev ^{aq}, C. Noeding ^{aa},

- A. Nomerotski ^{bc}, S.F. Novaes ^d, T. Nunnemann ^{ac}, E. Nurse ^{aw}, V. O'Dell ^{bc},
 D.C. O'Neil ^{e,f,g,h}, V. Oguri ^c, N. Oliveira ^c, N. Oshima ^{bc}, G.J. Otero y Garzón ^{bd},
 P. Padley ^{cd}, N. Parashar ^{bl}, S.K. Park ^{ai}, J. Parsons ^{bu}, R. Partridge ^{ca}, N. Parua ^{bw},
 A. Patwa ^{bx}, G. Pawloski ^{cd}, P.M. Perea ^{ba}, E. Perez ^u, P. Pétroff ^s, M. Petteni ^{av},
 R. Piegaia ^a, M.-A. Pleier ^{bv}, P.L.M. Podesta-Lerma ^{aj}, V.M. Podstavkov ^{bc},
 Y. Pogorelov ^{bh}, M.-E. Pol ^b, A. Pompoš ^{bz}, B.G. Pope ^{bq}, W.L. Prado da Silva ^c,
 H.B. Prosper ^{bb}, S. Protopopescu ^{bx}, J. Qian ^{bp}, A. Quadt ^z, B. Quinn ^{br}, K.J. Rani ^{ag},
 K. Ranjan ^{af}, P.A. Rapidis ^{bc}, P.N. Ratoff ^{au}, S. Reucroft ^{bo}, M. Rijssenbeek ^{bw},
 I. Ripp-Baudot ^{v,w}, F. Rizatdinova ^{bk}, S. Robinson ^{av}, R.F. Rodrigues ^c, C. Royon ^u,
 P. Rubinov ^{bc}, R. Ruchti ^{bh}, V.I. Rud ^{ao}, G. Sajot ^q, A. Sánchez-Hernández ^{aj},
 M.P. Sanders ^{bm}, A. Santoro ^c, G. Savage ^{bc}, L. Sawyer ^{bl}, T. Scanlon ^{av}, D. Schaile ^{ac},
 R.D. Schamberger ^{bw}, H. Schellman ^{bf}, P. Schieferdecker ^{ac}, C. Schmitt ^{ad},
 C. Schwanenberger ^z, A. Schwartzman ^{bt}, R. Schwienhorst ^{bq}, S. Sengupta ^{bb},
 H. Severini ^{bz}, E. Shabalina ^{bd}, M. Shamim ^{bk}, V. Shary ^u, A.A. Shchukin ^{ap},
 W.D. Shephard ^{bh}, R.K. Shivpuri ^{af}, D. Shpakov ^{bo}, R.A. Sidwell ^{bk}, V. Simak ^m,
 V. Sirotenko ^{bc}, P. Skubic ^{bz}, P. Slattery ^{bv}, R.P. Smith ^{bc}, K. Smolek ^m, G.R. Snow ^{bs},
 J. Snow ^{by}, S. Snyder ^{bx}, S. Söldner-Rembold ^{aw}, X. Song ^{be}, L. Sonnenschein ^t,
 A. Sopczak ^{au}, M. Sosebee ^{cb}, K. Soustruznik ^l, M. Souza ^b, B. Spurlock ^{cb},
 N.R. Stanton ^{bk}, J. Stark ^q, J. Steele ^{bl}, K. Stevenson ^{bg}, V. Stolin ^{an}, A. Stone ^{bd},
 D.A. Stoyanova ^{ap}, J. Strandberg ^{ar,as,at}, M.A. Strang ^{cb}, M. Strauss ^{bz}, R. Ströhmer ^{ac},
 D. Strom ^{bf}, M. Strovink ^{ay}, L. Stutte ^{bc}, S. Sumowidagdo ^{bb}, A. Sznajder ^c, M. Talby ^r,
 P. Tamburello ^{ax}, W. Taylor ^{e,f,g,h}, P. Telford ^{aw}, J. Temple ^{ax}, M. Tomoto ^{bc}, T. Toole ^{bm},
 J. Torborg ^{bh}, S. Towers ^{bw}, T. Trefzger ^{ab}, S. Trincaz-Duvold ^t, B. Tuchming ^u,
 C. Tully ^{bt}, A.S. Turcot ^{aw}, P.M. Tuts ^{bu}, L. Uvarov ^{aq}, S. Uvarov ^{aq}, S. Uzunyan ^{be},
 B. Vachon ^{e,f,g,h}, P.J. van den Berg ^{ak}, R. Van Kooten ^{bg}, W.M. van Leeuwen ^{ak},
 N. Varelas ^{bd}, E.W. Varnes ^{ax}, A. Vartapetian ^{cb}, I.A. Vasilyev ^{ap}, M. Vaupel ^{ad},
 P. Verdier ^x, L.S. Vertogradov ^{am}, M. Verzocchi ^{bm}, F. Villeneuve-Seguier ^{av},
 J.-R. Vlimant ^t, E. Von Toerne ^{bk}, M. Vreeswijk ^{ak}, T. Vu Anh ^s, H.D. Wahl ^{bb},
 L. Wang ^{bm}, J. Warchol ^{bh}, G. Watts ^{cf}, M. Wayne ^{bh}, M. Weber ^{bc}, H. Weerts ^{bq},
 N. Wermes ^z, A. White ^{cb}, V. White ^{bc}, D. Wicke ^{bc}, D.A. Wijngaarden ^{al}, G.W. Wilson ^{bj},
 S.J. Wimpenny ^{ba}, J. Wittlin ^{bn}, M. Wobisch ^{bc}, J. Womersley ^{bc}, D.R. Wood ^{bo},
 T.R. Wyatt ^{aw}, Q. Xu ^{bp}, N. Xuan ^{bh}, S. Yacoob ^{bf}, R. Yamada ^{bc}, M. Yan ^{bm}, T. Yasuda ^{bc},
 Y.A. Yatsunenko ^{am}, Y. Yen ^{ad}, K. Yip ^{bx}, H.D. Yoo ^{ca}, S.W. Youn ^{bf}, J. Yu ^{cb},
 A. Yurkewicz ^{bw}, A. Zabi ^s, A. Zatserklyaniy ^{be}, M. Zdrazil ^{bw}, C. Zeitnitz ^{ab}, D. Zhang ^{bc},
 X. Zhang ^{bz}, T. Zhao ^{cf}, Z. Zhao ^{bp}, B. Zhou ^{bp}, J. Zhu ^{bw}, M. Zielinski ^{bv}, D. Ziemińska ^{bg},
 A. Ziemiński ^{bg}, R. Zitoun ^{bw}, V. Zutshi ^{be}, E.G. Zverev ^{ao}

^a Universidad de Buenos Aires, Buenos Aires, Argentina^b LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

- ^c Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil
^d Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil
^e University of Alberta, Edmonton, AB, Canada
^f Simon Fraser University, Burnaby, BC, Canada
^g York University, Toronto, ON, Canada
^h McGill University, Montreal, PQ, Canada
ⁱ Institute of High Energy Physics, Beijing, People's Republic of China
^j University of Science and Technology of China, Hefei, People's Republic of China
^k Universidad de los Andes, Bogotá, Colombia
^l Center for Particle Physics, Charles University, Prague, Czech Republic
^m Czech Technical University, Prague, Czech Republic
ⁿ Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic
^o Universidad San Francisco de Quito, Quito, Ecuador
^p Laboratoire de Physique Corpusculaire, IN2P3-CNRS, Université Blaise Pascal, Clermont-Ferrand, France
^q Laboratoire de Physique Subatomique et de Cosmologie, IN2P3-CNRS, Université de Grenoble I, Grenoble, France
^r CPPM, IN2P3-CNRS, Université de la Méditerranée, Marseille, France
^s IN2P3-CNRS, Laboratoire de l'Accélérateur Linéaire, Orsay, France
^t LPNHE, IN2P3-CNRS, Universités Paris VI and VII, Paris, France
^u DAPNIA/Service de Physique des Particules, CEA, Saclay, France
^v IRéS, IN2P3-CNRS, Université Louis Pasteur, Strasbourg, France
^w Université de Haute Alsace, Mulhouse, France
^x Institut de Physique Nucléaire de Lyon, IN2P3-CNRS, Université Claude Bernard, Villeurbanne, France
^y III. Physikalisches Institut A, RWTH Aachen, Aachen, Germany
^z Physikalisches Institut, Universität Bonn, Bonn, Germany
^{aa} Physikalisches Institut, Universität Freiburg, Freiburg, Germany
^{ab} Institut für Physik, Universität Mainz, Mainz, Germany
^{ac} Ludwig-Maximilians-Universität München, München, Germany
^{ad} Fachbereich Physik, University of Wuppertal, Wuppertal, Germany
^{ae} Panjab University, Chandigarh, India
^{af} Delhi University, Delhi, India
^{ag} Tata Institute of Fundamental Research, Mumbai, India
^{ah} University College Dublin, Dublin, Ireland
^{ai} Korea Detector Laboratory, Korea University, Seoul, South Korea
^{aj} CINVESTAV, Mexico City, Mexico
^{ak} FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands
^{al} Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands
^{am} Joint Institute for Nuclear Research, Dubna, Russia
^{an} Institute for Theoretical and Experimental Physics, Moscow, Russia
^{ao} Moscow State University, Moscow, Russia
^{ap} Institute for High Energy Physics, Protvino, Russia
^{aq} Petersburg Nuclear Physics Institute, St. Petersburg, Russia
^{ar} Lund University, Lund, Sweden
^{as} Royal Institute of Technology and Stockholm University, Stockholm, Sweden
^{at} Uppsala University, Uppsala, Sweden
^{au} Lancaster University, Lancaster, United Kingdom
^{av} Imperial College, London, United Kingdom
^{aw} University of Manchester, Manchester, United Kingdom
^{ax} University of Arizona, Tucson, AZ 85721, USA
^{ay} Lawrence Berkeley National Laboratory and University of California, Berkeley, CA 94720, USA
^{az} California State University, Fresno, CA 93740, USA
^{ba} University of California, Riverside, CA 92521, USA
^{bb} Florida State University, Tallahassee, FL 32306, USA
^{bc} Fermi National Accelerator Laboratory, Batavia, IL 60510, USA
^{bd} University of Illinois at Chicago, Chicago, IL 60607, USA
^{be} Northern Illinois University, DeKalb, IL 60115, USA
^{bf} Northwestern University, Evanston, IL 60208, USA
^{bg} Indiana University, Bloomington, IN 47405, USA

- ^{bh} University of Notre Dame, Notre Dame, IN 46556, USA
- ^{bi} Iowa State University, Ames, IA 50011, USA
- ^{bj} University of Kansas, Lawrence, KS 66045, USA
- ^{bk} Kansas State University, Manhattan, KS 66506, USA
- ^{bl} Louisiana Tech University, Ruston, LA 71272, USA
- ^{bm} University of Maryland, College Park, MD 20742, USA
- ^{bn} Boston University, Boston, MA 02215, USA
- ^{bo} Northeastern University, Boston, MA 02115, USA
- ^{bp} University of Michigan, Ann Arbor, MI 48109, USA
- ^{bq} Michigan State University, East Lansing, MI 48824, USA
- ^{br} University of Mississippi, University, MS 38677, USA
- ^{bs} University of Nebraska, Lincoln, NE 68588, USA
- ^{bt} Princeton University, Princeton, NJ 08544, USA
- ^{bu} Columbia University, New York, NY 10027, USA
- ^{bv} University of Rochester, Rochester, NY 14627, USA
- ^{bw} State University of New York, Stony Brook, NY 11794, USA
- ^{bx} Brookhaven National Laboratory, Upton, NY 11973, USA
- ^{by} Langston University, Langston, OK 73050, USA
- ^{bz} University of Oklahoma, Norman, OK 73019, USA
- ^{ca} Brown University, Providence, RI 02912, USA
- ^{cb} University of Texas, Arlington, TX 76019, USA
- ^{cc} Southern Methodist University, Dallas, TX 75275, USA
- ^{cd} Rice University, Houston, TX 77005, USA
- ^{ce} University of Virginia, Charlottesville, VA 22901, USA
- ^{cf} University of Washington, Seattle, WA 98195, USA

Received 31 May 2005; received in revised form 25 July 2005; accepted 27 August 2005

Available online 15 September 2005

Editor: L. Rolandi

Abstract

We present a measurement of the top quark pair ($t\bar{t}$) production cross section ($\sigma_{t\bar{t}}$) in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using 230 pb $^{-1}$ of data collected by the DØ experiment at the Fermilab Tevatron Collider. We select events with one charged lepton (electron or muon), missing transverse energy, and jets in the final state. We employ lifetime-based b -jet identification techniques to further enhance the $t\bar{t}$ purity of the selected sample. For a top quark mass of 175 GeV, we measure $\sigma_{t\bar{t}} = 8.6^{+1.6}_{-1.5}(\text{stat. + syst.}) \pm 0.6(\text{lumi.})$ pb, in agreement with the standard model expectation.

© 2005 Published by Elsevier B.V.

PACS: 13.85.Lg; 13.85.Qk; 14.65.Ha

The top quark was discovered at the Fermilab Tevatron Collider by the CDF and DØ Collaborations [1] in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV based on about 50 pb $^{-1}$ of data per experiment. The increased statistics and higher collision energy of $\sqrt{s} = 1.96$ TeV of Tevatron Run II allow more precise measurements of top quark

properties, including its production and decay characteristics. Theoretical calculations performed within the framework of the standard model (SM) predict the $t\bar{t}$ production cross section ($\sigma_{t\bar{t}}$) with an uncertainty of less than 15% [2]. A significant deviation from this prediction would signal the presence of physics beyond the SM, such as $t\bar{t}$ resonant production [3]. The CDF and DØ Collaborations have previously reported measurements of $\sigma_{t\bar{t}}$ at $\sqrt{s} = 1.8$ TeV [4]. Recent measurements at $\sqrt{s} = 1.96$ TeV by the CDF [5] and

E-mail address: flera@fnal.gov (F. Rizatdinova).

¹ Visitor from University of Zurich, Zurich, Switzerland.

DØ [6] Collaborations agree with the SM prediction within their experimental uncertainties.

In the SM, the value of the Cabibbo–Kobayashi–Maskawa (CKM) matrix element $|V_{tb}|$ is tightly constrained by indirect measurements, $|V_{tb}| = 0.9990\text{--}0.9992$ (at 90% C.L.) [7]. If the CKM matrix is unitary and there are only 3 generations of leptons and quarks, the ratio of branching fractions $R = \mathcal{B}(t \rightarrow Wb)/\mathcal{B}(t \rightarrow Wq)$ can be interpreted as $|V_{tb}|^2$. A direct measurement of the ratio of branching fractions R [8] is found to be consistent with the SM prediction $R = 1$. In this analysis we assume that the top quark decays exclusively to a W boson and b -quark. The $t\bar{t}$ cross section was measured without this assumption in Ref. [6]. In the presented analysis, we look at the lepton + jets final state which results from the leptonic decay of one of the W bosons and the hadronic decay of the other. The event signature is one lepton with high transverse momentum, large transverse energy imbalance (\cancel{E}_T) due to the undetected neutrino, and four jets, two of which result from hadronization of the b quarks.

In this Letter, we report the measurement of $\sigma_{t\bar{t}}$ in the lepton (electron or muon) plus jets channel using b -jet identification (b -tagging) techniques exploiting the long lifetime of B hadrons. The data were collected by the DØ experiment from August 2002 through March 2004, and correspond to an integrated luminosity of $226 \pm 15 \text{ pb}^{-1}$ ($229 \pm 15 \text{ pb}^{-1}$) in the electron (muon) sample.

The DØ detector includes a tracking system, calorimeters, and a muon spectrometer [9]. The tracking system consists of a silicon microstrip tracker (SMT) and a central fiber tracker (CFT), both located inside a 2 T superconducting solenoid. The tracker design provides efficient charged particle measurements in the pseudorapidity region $|\eta| < 3$.² The SMT strip pitch of 50–80 μm allows a precise reconstruction of the primary interaction vertex (PV) and an accurate determination of the impact parameter of a track relative to the PV,³ which are the key components of the lifetime-

² Rapidity y and pseudorapidity η are defined as functions of the polar angle θ and parameter β as $y(\theta, \beta) \equiv \frac{1}{2} \ln[(1 + \beta \cos \theta)/(1 - \beta \cos \theta)]$ and $\eta(\theta) \equiv y(\theta, 1)$, where β is the ratio of a particle's momentum to its energy.

³ Impact parameter is defined as the distance of closest approach (d_{ca}) of the track to the primary vertex in the plane transverse to

based b -jet tagging algorithms. The PV is required to be within the SMT fiducial volume and consist of at least 3 tracks. The calorimeter consists of a central section (CC) covering $|\eta| < 1.1$, and two end calorimeters (EC) extending the coverage to $|\eta| \approx 4.2$. The muon system surrounds the calorimeter and consists of three layers of tracking detectors and two layers of scintillators [10]. A 1.8 T iron toroidal magnet is located outside the innermost layer of the muon detector. The luminosity is calculated from the rate for $p\bar{p}$ inelastic collisions detected using two hodoscopes of scintillation counters mounted close to the beam pipe on the front surfaces of the EC calorimeters.

We select data samples in the electron and muon channels by requiring an isolated electron with $p_T > 20 \text{ GeV}$ and $|\eta| < 1.1$, or an isolated muon with $p_T > 20 \text{ GeV}$ and $|\eta| < 2.0$. More details on the lepton identification as well as trigger requirements are reported elsewhere [6]. In both channels, we require \cancel{E}_T to exceed 20 GeV and not be collinear with the lepton direction in the transverse plane. These W boson candidate events must be accompanied by one or more jets with $p_T > 15 \text{ GeV}$ and rapidity $|y| < 2.5$ (footnote 2). Jets are defined using a cone algorithm with radius $\Delta R = 0.5$ [11]. We classify the selected events according to their jet multiplicity. Events with 3 or ≥ 4 jets are expected to be enriched in $t\bar{t}$ signal, whereas events with only 1 or 2 jets are expected to be dominated by background. We use the former to estimate $\sigma_{t\bar{t}}$, and the latter to verify the background normalization procedure.

The main background in this analysis is the production of W bosons in association with jets ($W + \text{jets}$), with the W boson decaying leptonically. In most cases, the jets accompanying the W boson originate from light (u, d, s) quarks and gluons ($W + \text{light jets}$). Depending on the jet multiplicity, between 2% and 14% of $W + \text{jets}$ events contain heavy flavor jets resulting from gluon splitting into $b\bar{b}$ or $c\bar{c}$ ($Wb\bar{b}$ or $Wc\bar{c}$, respectively), while in about 5% of events, a single c quark is present in the final state as a result of the W boson radiated from an s quark from the proton's or antiproton's sea (Wc). A sizeable background arises from strong production of two or more

the beamline. Impact parameter significance is defined as $d_{ca}/\sigma_{d_{ca}}$, where $\sigma_{d_{ca}}$ is the uncertainty on d_{ca} .

jets (“multijets”), with one of the jets misidentified as a lepton and accompanied by large \not{E}_T resulting from mismeasurements of jet energies. Significantly smaller contributions to the background arise from single top, $Z + \text{jets}$, and weak diboson (WW , WZ and ZZ) production. Only a small fraction of the background events contain b - or c -quark jets in the final state. As a consequence, the signal-to-background ratio is significantly enhanced when at least one jet is identified as a b -quark jet.

We use a secondary vertex tagging (SVT) algorithm to identify b -quark jets. Secondary vertices are reconstructed from two or more tracks satisfying the following requirements: $p_T > 1 \text{ GeV}$, ≥ 1 hits in the SMT layers and impact parameter significance $d_{ca}/\sigma_{d_{ca}} > 3.5$ (footnote 3). Tracks identified as arising from K_S^0 or Λ decays or from γ conversions are not considered. For a secondary vertex reconstructed within a jet, we define the decay length L_{xy} as the distance between primary and secondary vertices times the sign of the decay length projection onto the jet axis. If the secondary vertex has a decay length significance $L_{xy}/\sigma_{L_{xy}} > 7$,⁴ the jet is tagged as a b -quark jet. Events with exactly 1 (≥ 2) tagged jets are referred to as single-tag (double-tag) events. We treat single-tag and double-tag events separately because of their different signal-to-background ratios.

Secondary vertices with $L_{xy}/\sigma_{L_{xy}} < -7$ appear due to finite resolution of their characteristics after reconstruction, and define the “negative tagging rate”. The negative tagging rate is used to estimate the probability for misidentifying a light flavor (u, d, s quark or gluon) jet as a b -quark jet (the “mis-tagging rate”).

We estimate both the b -tagging efficiency and the mis-tagging rate using jets with ≥ 2 tracks satisfying less stringent requirements than those for SVT. In particular, the p_T cut is reduced from 1 to 0.5 GeV for all but the highest p_T track, and no cut on $d_{ca}/\sigma_{d_{ca}}$ of the tracks is made. These requirements have an efficiency per jet $> 80\%$ for $p_T > 30 \text{ GeV}$ and integrated over y . We measure the b -tagging efficiency in a data sample of dijet events with enhanced heavy fla-




Fig. 1. Measured b -tagging efficiency (circles) and mistagging rate (triangles), and estimated c -tagging efficiency (solid line) as a function of jet p_T .

vor content by requiring a jet with an associated muon at high transverse momentum relative to the jet axis. By comparing the SVT and muon-tagged jet samples, the tagging efficiency for semileptonic b -quark decays (“semileptonic b -tagging efficiency”) can be inferred. We make use of a Monte Carlo (MC) simulation to further correct the measured efficiency to the tagging efficiency for inclusive b -quark decays. We estimate the c -tagging efficiency from the same simulation, corrected by a scale factor defined as the ratio of the semileptonic b -tagging efficiency measured in data to that measured in the simulation. We estimate the mistagging rate from the negative tagging rate measured in dijet events, corrected for the contribution of heavy-flavor jets and the presence of long-lived particles in light-flavor jets. Fig. 1 shows the b -tagging efficiency, c -tagging efficiency and mis-tagging rate as a function of jet p_T .

We simulate $t\bar{t}$ production, and all background processes except multijets, using ALPGEN [12] to generate the parton-level processes, and PYTHIA [13] to provide fragmentation and to decay all unstable particles except B hadrons and τ leptons, which are modeled via EVTGEN [14] and TAUOLA [15], respectively. We process the generated events through the full GEANT-based [16] DØ detector simulation and the same reconstruction program used to process the data. We apply small additional smearing to the reconstructed objects to improve the agreement between the data and the simulation, and account for remaining discrepancies using correction factors derived by comparing the efficiencies measured in $Z \rightarrow \ell^+\ell^-$ data to

⁴ Decay length L_{xy} is defined as the distance from the primary to the secondary vertex in the plane transverse to the beamline. Decay length significance is defined as $L_{xy}/\sigma_{L_{xy}}$, where $\sigma_{L_{xy}}$ is the uncertainty on L_{xy} .

Table 1

Summary of observed and expected numbers of events before tagging and with exactly one jet tagged

	$W + 1$ jet	$W + 2$ jets	$W + 3$ jets	$W + \geq 4$ jets
<i>Before tagging</i>				
Observed	14054	5502	1365	367
Multijets	718 ± 78	516 ± 43	190 ± 14	66 ± 6
<i>After tagging</i>				
$W + \text{light}$	36.8 ± 4.0	21.4 ± 2.4	7.2 ± 0.9	1.8 ± 0.3
Wc	47.8 ± 5.4	24.2 ± 2.7	5.7 ± 0.7	0.8 ± 0.1
$Wc\bar{c}$	12.2 ± 3.4	17.2 ± 4.8	6.6 ± 1.9	2.2 ± 0.7
$Wb\bar{b}$	33.9 ± 8.7	43.2 ± 11.0	15.1 ± 3.9	4.5 ± 1.3
Multijets	14.9 ± 1.9	16.3 ± 2.1	8.3 ± 1.5	4.0 ± 1.2
Other bkg	6.6 ± 0.9	20.6 ± 2.1	8.2 ± 0.8	2.2 ± 0.4
Total bkg	152.4 ± 14.8	142.9 ± 16.0	51.0 ± 5.6	15.6 ± 1.7
$t\bar{t} \rightarrow \ell + \text{jets}$	0.4 ± 0.1	6.8 ± 1.4	24.4 ± 1.7	34.8 ± 4.3
Total expected	152.8 ± 14.8	149.7 ± 16.2	75.4 ± 5.9	50.4 ± 4.8
Observed	137	145	79	62

those obtained from the simulation. For all processes except the multijets background, we make use of the MC simulation to compute the total acceptance, applying the trigger, reconstruction and tagging efficiencies measured using data. The tagging probability for a particular process depends on the flavor composition of the jets in the final state as well as on the overall event kinematics. We estimate it by applying the tagging rates measured in data to each jet in the simulation, taking into consideration its flavor, p_T , and y . In the case of $W + \text{jets}$ events, we also use the simulation to estimate the fraction of the different $W + \text{heavy flavor}$ subprocesses.

We compute the $t\bar{t}$ acceptance for events with a true electron or muon arising from a $W \rightarrow \ell\nu$ ($\ell = e, \mu, \tau$) decay, corresponding to total branching fractions of 17.106% and 17.036% [7], respectively, in the electron and muon channels. In the electron channel, the total acceptance before tagging is estimated to be $(10.8 \pm 0.8)\%$ and $(14.2 \pm 1.7)\%$, for events with 3 and those with ≥ 4 jets, respectively. The corresponding numbers for the muon channel are $(9.9 \pm 1.0)\%$ and $(14.1 \pm 1.9)\%$. The estimated single-tag efficiencies are $(43.4 \pm 1.2)\%$ and $(45.3 \pm 1.0)\%$ for events with 3 and those with ≥ 4 jets, respectively. The corresponding double-tag efficiencies are $(10.4 \pm 1.0)\%$ and $(14.2 \pm 1.3)\%$.

We estimate the number of multijet events from the data for each jet multiplicity using the matrix method described in Ref. [6], separately for the samples before and after tagging. Smaller contributions from sin-

Table 2

Summary of observed and expected number of events with two or more jets tagged

	$W + 2$ jets	$W + 3$ jets	$W + \geq 4$ jets
$Wb\bar{b}$	5.7 ± 1.6	2.2 ± 0.6	0.7 ± 0.2
Other bkg	3.7 ± 0.4	2.0 ± 0.3	0.5 ± 0.3
Total bkg	9.4 ± 1.8	4.2 ± 0.8	1.2 ± 0.3
$t\bar{t} \rightarrow \ell + \text{jets}$	0.8 ± 0.2	5.9 ± 0.7	10.9 ± 1.9
Total expected	10.2 ± 1.9	10.1 ± 1.2	12.1 ± 2.0
Observed	15	9	14

gle top, $Z + \text{jets}$, and diboson production (collectively referred to as “other bkg”) are estimated from the simulation, normalized to the next-to-leading order theoretical cross sections [17,18]. We also include under “other bkg” the contribution from $t\bar{t}$ with both W bosons decaying leptonically, assuming the same $\sigma_{t\bar{t}}$ as for $t\bar{t} \rightarrow \ell + \text{jets}$. We determine the number of tagged $W + \text{jets}$ events as the product of the number of $W + \text{jets}$ events in data before tagging and the average tagging probability for $W + \text{jets}$ events (e.g., $\approx 4\%$ for single-tag and $\approx 0.4\%$ for double-tag events with ≥ 4 jets). The number of $W + \text{jets}$ events before tagging is computed as a difference between the number of selected events and the estimated contribution from the rest of processes (multijets, $t\bar{t}$, and “other bkg”).

Tables 1 and 2 summarize the sample composition for single-tag and double-tag events, respectively, assuming $\sigma_{t\bar{t}} = 7.0 \text{ pb}$. Fig. 2 shows the observed and expected number of events for each jet multiplicity.




Fig. 2. Expected and observed number of (a) single-tag and (b) double-tag events. The hatched area represents the total uncertainty in the expectation.

We interpret the excess over the background expectation in the third and fourth jet multiplicity bins as the $t\bar{t}$ signal. The good agreement between observation and expectation in the first and second jet multiplicity bins validates the background estimation procedure.

We calculate $\sigma_{t\bar{t}}$ by maximizing a likelihood function including a Poisson term for each of the eight independent channels considered: 3 and ≥ 4 jets, for single- and double-tag events in the electron and muon channels. At each step in the maximization, the multijet background in these eight tagged samples, and the corresponding samples before tagging, is constrained within errors to the amount determined by the matrix method. In addition, we include a Gaussian term for each of the systematic uncertainties considered, following the procedure described in Ref. [19]. In this approach, each source of systematic uncertainty is allowed to affect the central value of the cross section during the maximization procedure, thus yielding a combined statistical and systematic uncertainty on $\sigma_{t\bar{t}}$. Assuming a top quark mass (m_t) of 175 GeV, we measure

$$\sigma_{t\bar{t}} = 8.6^{+1.6}_{-1.5}(\text{stat. + syst.}) \pm 0.6(\text{lumi.}) \text{ pb},$$

in good agreement with the SM prediction of 6.77 ± 0.42 pb [2].

The contribution due to each individual source of systematic uncertainty can be estimated by redoing the fit after fixing all but the corresponding Gaussian term and unfolding the statistical uncertainty from the resulting total uncertainty. The statistical uncertainty of $+1.2_{-1.1}$ pb is obtained from the fit where all Gaussian terms are fixed. As shown in Table 3, b -jet tagging

Table 3
Summary of systematic uncertainties on $\sigma_{t\bar{t}}$

Source	$\Delta\sigma_{t\bar{t}}$ (pb)
b -tagging efficiency	$+0.6$ -0.5
Jet energy calibration	$+0.5$ -0.4
Background modeling	± 0.5
Lepton selections	$+0.5$ -0.4
Jet identification	$+0.3$ -0.2
Multijet background	$+0.3$ -0.2
Mis-tagging rate	± 0.1
Total	$+1.1$ -1.0

efficiency, jet energy calibration [6], and background modeling are the leading sources of systematic uncertainty. In addition, a systematic uncertainty of 6.5% from the luminosity measurement [20] has been assigned. In the top quark mass range of 160 to 190 GeV, the measured cross section decreases (increases) by 0.06 pb per 1 GeV shift of m_t above (below) 175 GeV.

We used an alternative b -tagging algorithm to cross check this result. This algorithm relies on counting tracks with significant impact parameter (CSIP) with respect to the PV: a jet is tagged if ≥ 2 (≥ 3) associated tracks have $d_{ca}/\sigma_{d_{ca}} > 3$ ($d_{ca}/\sigma_{d_{ca}} > 2$). As compared to SVT, this algorithm has a slightly higher b -tagging efficiency and about a factor of two higher mis-tagging rate. The measured cross section using the CSIP algorithm is $\sigma_{t\bar{t}} = 7.6^{+1.7}_{-1.4}(\text{stat. + syst.}) \pm 0.5(\text{lumi.})$ pb, consistent with the SVT result once the existing overlap between both samples is taken into account. While we are currently not combining these two results, the

fact that different b -tagging techniques are only partially correlated will be exploited in future analyses to further increase the precision of this measurement.

In summary, we have measured the $t\bar{t}$ production cross section in $p\bar{p}$ interactions at $\sqrt{s} = 1.96$ TeV in the lepton + jets channel using lifetime b -tagging. Our measurement yields $\sigma_{t\bar{t}} = 8.6^{+1.6}_{-1.5}$ (stat. + syst.) ± 0.6 (lumi.) pb, in a good agreement with the SM prediction.

Acknowledgements

We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); FASI, Rosatom and RFBR (Russia); CAPES, CNPq, FAPERJ, FAPESP and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF (Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); PPARC (United Kingdom); MSMT (Czech Republic); CRC Program, CFI, NSERC and WestGrid Project (Canada); BMBF and DFG (Germany); SFI (Ireland); Research Corporation, Alexander von Humboldt Foundation, and the Marie Curie Program.

References

- [1] CDF Collaboration, F. Abe, et al., Phys. Rev. Lett. 74 (1995) 2626;
DØ Collaboration, S. Abachi, et al., Phys. Rev. Lett. 74 (1995) 2632.
- [2] R. Bonciani, et al., Nucl. Phys. B 529 (1998) 424;
M. Cacciari, et al., JHEP 0404 (2004) 068;
N. Kidonakis, R. Vogt, Phys. Rev. D 68 (2003) 114014.
- [3] C.T. Hill, S.J. Parke, Phys. Rev. D 49 (1994) 4454;
H.P. Nilles, Phys. Rep. 110 (1984) 1;
H.E. Haber, G.L. Kane, Phys. Rep. 117 (1985) 75.
- [4] CDF Collaboration, T. Affolder, et al., Phys. Rev. D 64 (2001) 032002;
DØ Collaboration, V. Abazov, et al., Phys. Rev. D 67 (2003) 012004.
- [5] CDF Collaboration, D. Acosta, et al., Phys. Rev. Lett. 93 (2004) 142001;
CDF Collaboration, D. Acosta, et al., Phys. Rev. D 71 (2005) 072005;
CDF Collaboration, D. Acosta, et al., Phys. Rev. D 71 (2005) 052003.
- [6] DØ Collaboration, V. Abazov, et al., Phys. Lett. B 626 (2005) 45, the next Letter in this issue.
- [7] S. Eidelman, et al., Phys. Lett. B 592 (2004) 1.
- [8] CDF Collaboration, T. Affolder, et al., Phys. Rev. Lett. 86 (2001) 3233.
- [9] DØ Collaboration, V. Abazov, et al., The upgraded DØ detector, in preparation for submission to Nucl. Instrum. Methods Phys. Res. A.
- [10] V. Abazov, et al., FERMILAB-PUB-05-034-E (2005).
- [11] We use the iterative, seed-based cone algorithm including midpoints, as described in G.C. Blazey, et al., in: U. Baur, R.K. Ellis (Eds.), Proceedings of the Workshop QCD and Weak Boson Physics in Run II, p. 47;
D. Zeppenfeld, FERMILAB-PUB-00-297 (2000).
- [12] M.L. Mangano, et al., JHEP 0307 (2003) 001.
- [13] T. Sjöstrand, et al., Comput. Phys. Commun. 135 (2001) 238.
- [14] D. Lange, Nucl. Instrum. Methods Phys. Res. A 462 (2001) 152.
- [15] S. Jadach, et al., Comput. Phys. Commun. 76 (1993) 361.
- [16] R. Brun, F. Carminati, CERN Programming Library Long Writeup W5013 (1993).
- [17] M.C. Smith, S. Willenbrock, Phys. Rev. D 54 (1996) 6696;
T. Stelzer, et al., Phys. Rev. D 56 (1997) 5919.
- [18] J.M. Campbell, R.K. Ellis, Phys. Rev. D 60 (1999) 113006.
- [19] P. Sinervo, in: L. Lyons, R.P. Mount, R. Reitmeyer (Eds.), Proceedings of Statistical Methods in Particle Physics, Astrophysics, and Cosmology, SLAC, Stanford, 2003, p. 334.
- [20] T. Edwards, et al., FERMILAB-TM-2278-E (2004).