



# B hadron rare decays and lifetimes

$B_s \rightarrow \mu\mu$  and  $B_d \rightarrow \mu\mu$   
B meson and  $\Lambda_b$  lifetimes



Marj Corcoran for the CDF and D0 collaborations


# The TeVatron is working well!



Many measurements are now within reach for the first time!


# $B_s \rightarrow \mu\mu$ and $B_d \rightarrow \mu\mu$

$B_s \rightarrow \mu\mu$  and  $B_d \rightarrow \mu\mu$  are FCNC in the Standard Model and are therefore highly suppressed.  $B_d \rightarrow \mu\mu$  is suppressed even further by  $V_{td}/V_{ts}$ .



Standard Model expectations :  
 $BR(B_s \rightarrow \mu\mu) = (3.42 \pm 0.54) \times 10^{-9}$   
 $BR(B_d \rightarrow \mu\mu) = (1.0 \pm 0.14) \times 10^{-10}$

**Many models beyond the SM enhance these BRs by orders of magnitude**  
 And different models can effect  $B_d$  and  $B_s$  differently.



$B_s \rightarrow \mu\mu$  at D0 –first result using L0!



L0 goes into the heart of D0 in the summer of 2006 with less than 1mm clearance!

RunIIa= pre-upgrade and RunIIb=post-upgrade.



# $B_s \rightarrow \mu\mu$ using $2 \text{ fb}^{-1}$

Data selection:

Two muons with  $p_t > 2.5 \text{ GeV}$  and  $|\eta| < 2$  which form a good vertex

Dimuon mass between 4.5 and 7 GeV

Dimuon  $p_t > 5 \text{ GeV}$ .

For RunIIa, there are 163K candidates

For RunIIb, there are 36K candidates

Early RunIIb data had several problems (firmware, timing).

Some are recoverable, some not, all problems are now fixed.

Background distributions are taken from the sidebands in  $M_{\mu\mu}$ .

Signal distributions are taken from Monte Carlo.



# $B_s \rightarrow \mu\mu$ using $2 \text{ fb}^{-1}$

Form a likelihood ratio based on six variables:

Muon pair isolation ( $\Delta R=1$ )


Decay length significance

Pointing angle  $\alpha$  (see below)

B impact parameter


Minimum muon impact parameter

Vertex  $\chi^2$  probability







# $B_s \rightarrow \mu\mu$ using $2 \text{ fb}^{-1}$



The effects of L0 are apparent in the improved uncertainty in the transverse decay length. “p17” is the pre-L0 data; “p20” is the post-L0 data.




$$B_s \rightarrow \mu\mu$$



Likelihood ratios for signal MC and background taken from the side bands in dimuon mass. The red arrow shows the optimized cut location.


# $B_s \rightarrow \mu\mu$ at D0

DØ Run IIa Preliminary




For the pre-upgrade data, there is one event in the signal region in mass and likelihood ratio. Expected background is  $0.8 \pm 0.2$  events.


DØ Run IIb Preliminary



For the post-upgrade data, there are two events in the signal region, with an expected background of  $1.1 \pm 0.3$  events.



$$B_s \rightarrow \mu\mu$$



Normalization mode is  
 $B^+ \rightarrow J/\psi K^+$  with  $J/\psi \rightarrow \mu\mu$

### Preliminary results:

From RunIIa:  $\text{BR}(B_s \rightarrow \mu\mu) < 9.5 \times 10^{-8}$  at 95% CL

From RunIIb:  $\text{BR}(B_s \rightarrow \mu\mu) < 4.0 \times 10^{-7}$  at 95% CL

Combined 95% CL limit is  $9.3 \times 10^{-8}$

Recall that the SM expectation is  $3.42 \times 10^9$



# $B_s \rightarrow \mu\mu$ and $B_d \rightarrow \mu\mu$ at CDF

CDF's results are based on 780 pb<sup>1</sup> of data. They have limits for both  $B_s$  and  $B_d$ .

Data selection:


The data are divided into two subsets-- one subset with both muons in the central muon system (CMU-CMU), and the second subset with one muon in the central muon extension (CMU-CMX).

Muon  $p_t > 2$  GeV and  $|\eta| < 0.6$  in CMU

Muon  $p_t > 2.2$  GeV and  $0.6 < |\eta| < 1.0$  in CMX



# $B_s \rightarrow \mu\mu$ and $B_d \rightarrow \mu\mu$ at CDF




A likelihood ratio is formed based on three variables: Isolation of the B candidate, 3D pointing angle, and  $P(\hat{\lambda}) = \exp(-\lambda/c\tau_A)$  where  $\lambda=c\tau$  for a given event and  $\tau_A$  is the world average lifetime.

Background distributions are taken from sidebands in  $M_{\mu\mu}$ . Signal distributions are from Monte Carlo.





# $B_s \rightarrow \mu\mu$ and $B_d \rightarrow \mu\mu$ at CDF



Normalization mode is  
 $B^+ \rightarrow J/\psi K$  with  $J/\psi \rightarrow \mu\mu$



# $B_s \rightarrow \mu\mu$ and $B_d \rightarrow \mu\mu$ at CDF



Likelihood ratio vs  $M_{\mu\mu}$ . One event is observed for  $B_s \rightarrow \mu\mu$  and two events are observed for  $B_d \rightarrow \mu\mu$ . Expected backgrounds are  $1.27 \pm 0.37$  for  $B_s$  and  $2.45 \pm 0.40$  for  $B_d$ .



# $B_s \rightarrow \mu\mu$ and $B_d \rightarrow \mu\mu$ at CDF

**Preliminary** CDF result for these decays:

$$\text{BR}(B_s \rightarrow \mu\mu) < 1.0 \times 10^{-7} \text{ at } 95\% \text{ CL}$$

$$\text{BR}(B_d \rightarrow \mu\mu) < 2.3 \times 10^{-8} \text{ at } 95\% \text{ CL.}$$

Recall the SM expectation

$$\text{BR}(B_s \rightarrow \mu\mu) = 3.42 \times 10^{-9}$$

$$\text{BR}(B_d \rightarrow \mu\mu) = 1.0 \times 10^{-10}$$

# B hadron lifetime measurements

B hadron lifetimes can be calculated in Heavy Quark Expansion models. To lowest order all b hadron lifetimes would be the same. To second and third order splittings occur but are expected to be small.

Expect  $\tau(B^+) > \tau(B^0) \sim \tau(B_s) > \tau(\Lambda_b)$ .

D0 has recent measurements:

$\Lambda_b \rightarrow J/\psi \Lambda$  with  $J/\psi \rightarrow \mu\mu$

$\Lambda_b \rightarrow \Lambda_c \mu \nu X$  (semileptonic)

CDF has recent measurements:


$\Lambda_b \rightarrow J/\psi \Lambda$  with  $J/\psi \rightarrow \mu\mu$

Precision lifetime measurements of  $B^+$ ,  $B^0$ , and  $B_s$ .




# B meson lifetime measurements from CDF

Care in understanding resolution functions and backgrounds yield precision measurements of the B meson lifetimes in the channels, based on  $1\text{fb}^1$  of data.



$B^+$  mass showing the sidebands used to estimate the background.




A maximum likelihood fit on three variables (mass,  $\tau$ , and error on  $\tau$ ) yields the lifetime




# $\Lambda_b$ lifetime measurement from CDF

Using the same method as in the high-statistics modes, CDF has measured the  $\Lambda_b$  lifetime in the channel  $\Lambda_b \rightarrow J/\psi \Lambda$  with  $J/\psi \rightarrow \mu\mu$  and  $\Lambda \rightarrow p\pi$ , based on  $1\text{fb}^{-1}$  of data.



Reconstructed  $\Lambda_b$  mass




A maximum likelihood fit on three variables (mass,  $\tau$ , and error on  $\tau$ ) yields the lifetime.




# $\Lambda_b$ lifetime measurements from D0


Based on  $1\text{fb}^{-1}$ , the  $\Lambda_b$  is observed in the decay  $\Lambda_b \rightarrow J/\psi \Lambda$  with  $J/\psi \rightarrow \mu\mu$  and  $\Lambda \rightarrow p\pi$ . A  $B^0$  lifetime measurement is also made with this dataset.



$\Lambda_b$  mass,  $174 \pm 21$  events




Maximum likelihood proper decay time fit



# $\Lambda_b$ lifetime measurement in the semileptonic channel

D0 has also made a measurement of the  $\Lambda_b$  lifetime in the semileptonic channel  $\Lambda_b \rightarrow \Lambda_c^+ \mu^- \bar{\nu} X$  with  $\Lambda_c \rightarrow p K_s$



Corrections must be made to the observed  $p$  for the missing neutrino, based on Monte Carlo.

$\chi^2$  fits to the lifetime are done in bins of visible proper decay length.


Example  $\Lambda_c$  mass plot for one bin of visible decay length.



Recent D0 and CDF measurements are **preliminary**.



The world average does not include the most recent measurements.

There appears to be a discrepancy of about 3.2 sigma in  $\Lambda_b$  lifetime measurements from the world average and the most recent CDF measurements.




Thanks to CDF for preparing this and other comparison plots!

# B hadron lifetime measurements



Other measurements are in good agreement.

# B hadron lifetime measurements



The most sensitive comparison to calculations is the ratio, which is in good agreement for B mesons.

Expectation from heavy quark expansion for  $\tau(B^+)/\tau(B^0)=1.04-1.08$ , in good agreement with observation.

# Conclusions

## **Spectacular B physics is being done at CDF and D0!**

Limits on  $B_s \rightarrow \mu\mu$  are still well above the SM expectation, but they are starting to push on expectations from physics beyond the SM.

$$\text{BR}(B_s \rightarrow \mu\mu) < 1.0 \times 10^{-7} \text{ (CDF)}$$

$$\text{BR}(B_d \rightarrow \mu\mu) < 2.3 \times 10^{-8} \text{ (CDF)}$$

$$\text{BR}(B_s \rightarrow \mu\mu) < 9.3 \times 10^{-8} \text{ (D0)}$$

Precision lifetime measurements are being made of B mesons and hadrons. There is perhaps a  $3\sigma$  disagreement in  $\Lambda_b$  lifetime, otherwise all measurements are consistent.

# Prospects

These results are based on at most  $2\text{fb}^1$  of data; many on  $1\text{fb}^1$  or less. Both experiments now have more than  $2\text{fb}^1$  recorded, and the data continues to come in at an impressive rate.

There are many results which I could not show, in particular CDF rare decays. Visit the CDF and D0 web sites!

[www-cdf.fnal.gov/physics/new/bottom/bottom.html](http://www-cdf.fnal.gov/physics/new/bottom/bottom.html)

[www-d0.fnal.gov/Run2Physics/WWW/results/b.htm](http://www-d0.fnal.gov/Run2Physics/WWW/results/b.htm)

Thanks to CDF B group for their help!  
Thanks to my D0 colleagues who did all the work.

