Single top quark physics at the Tevatron and the LHC

Reinhard Schwienhorst

MICHIGAN STATE UNIVERSITY

Wuppertal seminar, June 2011

Outline

- Introduction
- Tevatron
  - SM
  - New physics
- LHC
- Conclusions


Higgs
boson
Electroweak
symmetry breaking
W boson


top quark
Coupling
strength ~ 1
Higgs
boson
Electroweak
symmetry breaking
W boson


July 2010
LEP2 and Tevatron (prel.)
LEP1 and SLD
68% CL
80.5
80.4
80.3
m_W [GeV]
m_H [GeV]
114
300
1000
$\Delta\alpha$
150
175
200
m_t [GeV]


top quark
Higgs boson
W boson

Key to electroweak symmetry breaking



Top quark decay



t-channel single top quark production





Wt associated production





s-channel single top production





SM single top quark production









SM tasks:

- measure total cross section → CKM matrix element $|V_{tb}|$
- top polarization, PDFs

New physics in single top

New interactions: FCNC



New coupling: modified Wtb coupling or anomalous CKM matrix



New particles: heavy boson or charged Higgs or T′ or B′



New physics tasks:

- measure individual cross sections → different modes sensitive to different new physics
- look for specific new physics scenarios
  - W’, fourth generation quarks, charged Higgs
  - gluon FCNC
  - anomalous Wtb couplings

Experimental setup: Fermilab Tevatron in Run II



Proton-antiproton collider

CM energy 1.96 TeV

→ ***Energy frontier - until a year ago***

Instantaneous luminosity >4E32 $cm^{-2}s^{-1}$

– >5 interactions per crossing, 1.7M crossing per second

→ ***Proton-antiproton luminosity frontier***

Fermilab single top history

Publication history

	· Search:	PRD 63, 031101 (2000)
	· Search:	PLB 517, 282 (2001)
Run I	· Search:	PLB 622, 265 (2005)
	· W':	PLB 641, 423 (2006)
	· Search:	PRD 75, 092007 (2007)
Run II	· Evidence:	PRL 98, 181802 (2007)
	· FCNC:	PRL 99, 191802 (2007)
	· W':	PRL 100, 211802 (2007)
	· Evidence:	PRD 78, 012005 (2008)
	· Wtb:	PRL 101, 221801 (2008)
	· Wtb:	PRL 102, 092002 (2009)
	· H^+:	(PRL) arXiv:0807.0859
	· Observation:	(PRL) arXiv:0903.0850
	. tau search	PLB 690, 5 (2010)
	. t-chan meas.	PLB 682, 363 (2010)
	. FCNC	PLB 693, 81 (2010)
	. top width	PRL 106, 22001 (2011)
	. W'	PLB 699, 145 (2011)
	. t-chan obs.	(PLB) arXiv:1105.2788

	· Search:	PRD 65, 091102 (2002)
	· W'	PRL 90, 081802 (2003)
	· Search:	PRD 69, 052003 (2004)
	· Search:	PRD 71, 012005 (2005)
	· Evidence:	PRL 101, 252001 (2008)
	· FCNC:	(PRL) arXiv:0812.3400
	· W':	(PRL) arXiv:0902.3276
	· Observation:	(PRL) arXiv:0903.0885
	. MET+jets	PRD 81, 072003 (2010)
	. Observation	PRD 82, 112005 (2010)

Single top quark event signature



Single top quark event signature



s-channel



t-channel





Background processes





Discriminating variables



cross-check sample







W+Jets Cross-Check Sample





Discriminating variables



Object kinematics

Event kinematics

Angular correlations

Jet reconstruction

Top reconstruction

- Started from ~ 600 variables
- Considered ~200 for multivariate filters
- Chose 97 depending on method and channel

Tevatron analysis outline

Trigger selection



Single top event kinematics



b-quark tagging





Multivariate techniques

BDT



NN



ME



Combination



Statistical analysis



Multivariate methods

Input:
discriminating variables

Method:
multivariate analysis

Output:
signal likelihood



Cut-Based



Neural networks



Decision trees



Likelihood



Boosted decision trees



Bayesian neural networks



Neuro evolution



Matrix Elements



Single top production cross section

- Several multivariate filters combined in one final discriminant
  - BNN (DØ)
  - NEAT (CDF)

Boosted Decision Trees | **Neural Networks** | **Matrix Elements** | **Likelihood**







Cross sections









Tevatron combination

- Each experiment observes 5 σ significance
- DØ and CDF measurements are consistent to 1.6 σ
- Combination of MVA distributions
  - Bayesian statistical analysis
  - Taking all uncertainty correlations into account







CKM matrix element Vtb





$$V_{CKM} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix}$$

- Cross section ∝ $|V_{tb}|^2$
  - Assume SM top quark decay: $|V_{td}|^2+|V_{ts}|^2 \ll |V_{tb}|^2$
  - Pure V-A and CP conserving Wtb vertex
- No assumption on number of families or unitarity



t-channel production

- New analysis with 5.4 fb^{-1}
- Dedicated t-channel filter
- No assumption about s-channel





- 5.5 σ significance







Top quark properties

- top quark width from t-channel cross section
  - use R=B(t→Wb)/B(t→Wx) to obtain full width



$$\Gamma_t = 2.05^{+0.57}_{-0.52}\ \text{GeV} \qquad \tau_t = (3.2^{+1.1}_{-0.7}) \times 10^{-25}\ \text{s}$$

- Top quark polarization from spin correlation





- Analysis using likelihood function
- Test for right-handed coupling in top production

$$polarization = \frac{\sigma_R - \sigma_L}{\sigma_R + \sigma_L} = -1^{+1.5}_{-0}$$

Tevatron new physics searches

Gluon-FCNC in single top



- CDF: 2→1 process
  - ★multivariate analysis using NN



$$\frac{\kappa_{gtu}}{\Lambda} < 0.025\ \mathrm{TeV}^{-1}$$

$$\frac{\kappa_{gtc}}{\Lambda} < 0.105\ \mathrm{TeV}^{-1}$$



- D0: 2→2 process
  - ★Same final state as t-channel
  - ★multivariate analysis using BNN

95% CL limits:	tgu	tgc
Cross section	0.20 pb	0.27 pb
κ_{tgf}/Λ	0.013 TeV^{-1}	0.057 TeV^{-1}
$\mathcal{B}(t \to qg)$	2.0×10^{-4}	3.9×10^{-3}

Single top anomalous coupling

- Left-vector (f^L_1, =1 in SM), right-vector (f^R_1), left-tensor (f^L_2), right-tensor (f^R_2)



$$\mathcal{L} = -\frac{g}{\sqrt{2}}\bar{b}\gamma^\mu V_{tb}(f_1^L P_L + f_1^R P_R)tW_\mu^- - \frac{g}{\sqrt{2}}\bar{b}\frac{i\sigma^{\mu\nu}q_\nu}{M_W}(f_2^L P_L + f_2^R P_R)tW_\mu^- + h.c.$$

- Single top is sensitive to magnitude (PRL 101, 221801 (2008))
- ttbar to ratios of couplings (W helicity, PRL 100, 062004 (2008))
- Best sensitivity through combination (PRL 102, 092002 (2009))



Search for W' boson



- Explore simultaneously left-handed and right-handed couplings
- 95% CL limits:

$M_{W'} > 863$ GeV (L only)

$M_{W'} > 885$ GeV (R, $M_{W'} < m_{nR}$)

$M_{W'} > 890$ GeV (R, $M_{W'} > m_{nR}$)

$M_{W'} > 916$ GeV (L+R)







Experimental setup: LHC at CERN

Proton-proton collider

CM energy 7 TeV

→ *Energy frontier*

Instantaneous luminosity > 6E32 $cm^{-2}s^{-1}$

– >8 interactions per crossing, 1.7M crossing per second

→ *Luminosity frontier*

Measurement of t-channel production



- 36 pb^{-1} of 2010 data analysis
- 156 pb^{-1} of 2011 ATLAS data analysis shown here



Event selection:

- 1 electron or muon E_T (p_T) > 25 GeV
- E_T^{Miss} > 25 GeV
- = 2 jets (p_T > 25 GeV), = 1 b-tag
- M_T > 60 GeV - E_T^{miss} (multijet veto)
- fewer channels than Tevatron, tighter cuts

Background Modeling









Cut-based t-channel analysis

- | untagged jet η | > 2.0
- 140 GeV < top quark mass < 190 GeV
- $|\Delta\eta$(lepton, b-tagged jet)| < 1.5
- | b-tagged jet η | < 2.0
- H_T > 180 GeV
- separate by lepton flavor and top charge





Neural network result



- 22 input variables





t-channel result

- Simple event-counting
- Expected cross section systematic uncertainties
  - B-tagging ~ 12%
  - MC statistics ~ 11%
  - ISR/FSR ~ 6%
  - Signal modeling $^{+36}_{-20}$%
  - Total systematics $^{+55}_{-29}$%
- Statistical uncertainty 17%

Cut-based: Observed cross section:
$\sigma_t = 97\ ^{+54}_{-30}$ pb
Observed (expected) significance:
6.3 (4.5) σ



NN: Observed cross section:
$\sigma_t = 76\ ^{+41}_{-21}$ pb

Observed (expected) significance:
6.2 (5.7) σ

t-channel result

- 2010 data, 2 analyses: 37-variable BDT and fit to spin correlation and light quark jet η







combined: observed cross section:
$\sigma_t = 83.6 \pm 29.8$ pb

Observed significance: 3.5 σ
Observed $|V_{tb}| > 0.68$ at 95% CL

- submitted to PRL

t-channel summary



Wt associated production



- Never before seen!
- search in lepton+jets and di-leptons



Dilepton backgrounds:

Top quark pairs

- Model using MC@NLO
- Normalize to data



Drell-Yan

- Model using Alpgen
- Normalize to data



- set limit of 158 pb (10 × σ_{Wt}^{SM})



Smaller backgrounds from $Z \rightarrow \tau\tau$, dibosons, W+jets, multijets

LHC new physics searches

ATLAS FCNC search



- first single top new physics search
- single-top production









ATLAS FCNC search



- first single top new physics search
- single-top production





First of many new physics searches!



Summary/Outlook

- Single top production observed at Tevatron and LHC
  - -Tevatron s-channel+t-channel observation in 2009
  - -Tevatron t-channel isolation in 2011
  - -LHC t-channel observation in 2011
  - -LHC first limit on Wt
  - -Next: Wt and s-channel observation

- Single top as a tool to look for new physics
  - -Tevatron: W′, FCNC, H+, anomalous couplings
  - -LHC: FCNC
  - -Next: many more LHC new physics searches

Larger LHC datasets will bring separate observation of all single top processes and many new physics searches