Single Top Quark Physics at DØ in Run II

Reinhard Schwienhorst

University of Wisconsin High Energy Seminar, 10/10/2005

Outline

- Introduction
  - Electroweak top quark production
- Search for single top at DØ
  - DØ detector
  - Neural Networks
  - Likelihoods
  - Results
- Future studies of top quark couplings
- Conclusions

Fundamental Fermions



Fundamental Fermions



King of the Fermions





Electroweak Symmetry Breaking



Yukawa coupling to Higgs field v

$$m_t = y_t\, v$$

$$y_t \sim 1$$







SM Higgs boson mass estimate





Key to electroweak symmetry breaking

Focus of many studies at the Tevatron

Top Quark Electroweak Charged Current Interaction



top quark decay

Electroweak Production of Top at the Tevatron



Electroweak Production of Top at the Tevatron

s-channel



$O(\alpha_s)$ corrections:



Electroweak Production of Top at the Tevatron

s-channel





Cao, RS, Yuan PRD74, 054023 (2005)

Electroweak Production of Top at the Tevatron

t-channel



Electroweak Production of Top at the Tevatron

t-channel



$O(\alpha_s)$ corrections:



Electroweak Production of Top at the Tevatron

t-channel





Cao, RS, Benitez, Brock, Yuan hep-ph/0504230



Tevatron Single Top Goals

- Observe single top quark production

Tevatron Single Top Goals

- Observe single top quark production
- Measure production cross sections $\rightarrow V_{tb}$



CKM Matrix

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

Tevatron Single Top Goals

- Observe single top quark production
- Measure production cross sections $\rightarrow V_{tb}$
- Look for physics beyond the Standard Model
  - Different sensitivity for s-channel and t-channel

Example:Top-Flavor





Tevatron Single Top Goals

- Observe single top quark production
- Measure production cross sections $\rightarrow V_{tb}$
- Look for physics beyond the Standard Model
- Study top quark spin correlations – probe V-A
  - Physics with ~100% polarized top quarks

Tevatron Single Top Goals

- Observe single top quark production
- Measure production cross sections $\rightarrow V_{tb}$
- Look for physics beyond the Standard Model
- Study top quark spin correlations – probe V-A
- Irreducible background in searches for associated Higgs production



Gateway to the Higgs

Tevatron Single Top Status

s-channel



t-channel



- Production cross section:

	s-channel	*t-channel*	*s+t*
– NLO calculation:	**0.88pb (±8%)**	**1.98pb (±11%)**	
– Run I 95% CL limits, DØ:	< 17pb	< 22pb	
CDF:	< 18pb	< 13pb	< 14pb
– Run II CDF 95% CL limits:	< 14pb	< 10pb	< 18pb

- Other Standard Model production mode (*Wt*) negligible


SPS
Site de Prevessin
SPS2
BA2
SPS3
BA3
SPS4
BA4
SPS1
BA1
FRANCE
SUISSE
SPS7
BA7
SPS6
BA6
SPS5
BA5
Site de Meyrin

WHERE IN THE
WORLD



Single Top

HERA
PETRA

Single Top at LEP and Hera: FCNC

- LEP:
  - $e^+e^- \rightarrow tc$

- Hera:
  - $ep \rightarrow et$







Experimental Detection of Single Top Events

Experimental Setup: Fermilab Tevatron in Run II

Proton-Antiproton Collider
CM Energy 1.96TeV
→ ***Energy Frontier***
Instantaneous luminosity 140E30
→ ***Luminosity Frontier***

Experimenters

- 19 countries
- 80 institutions
- 670 physicists

Apparatus: Run II DØ Detector



Analysis Outline

Goal:

Maximize Sensitivity







1. Event Selection
   - Select *W*-like events
   - Maximize acceptance
   - Model backgrounds
2. Event Analysis
   - Find discriminating variables
   - Cut/combine in multivariate analysis
3. Determine cross section
   - Event counting or Binned likelihood
   - Separately for s-channel, t-channel, and combined

Event Selection

- Trigger:
  - Electron + ≥1 jets, muon + ≥1 jets
- Lepton:
  - 1 electron: $p_T > 15\text{GeV}$, $|\eta^{det}|<1.1$
  - 1 muon: $p_T > 15\text{GeV}$, $|\eta^{det}|<2.0$
- Neutrino: $\not{E}_T > 15\text{GeV}$
- Jets:
  - $p_T > 15\text{GeV}$, $|\eta^{det}|<3.4$, p_T (jet 1)> 25GeV, $|\eta^{det}(\text{jet1})|<2.5$
  - $2 \leq n_{jets} \leq 4$
- Reject mis-reconstructed events





Event Selection: b-tagging

Algorithms:
- Secondary vertex tag
- Impact parameter (jet lifetime) tag



s-channel

- Final state:
  - 2 high-p_T b-jets
- ➔ Require ≥1 b-tagged jet



t-channel

- Final state:
  - 1 high-p_T b-jet
  - 1 high-p_T light quark jet
- ➔ Require ≥1 b-tagged jet
- ➔ Require ≥1 untagged jet

Background Modeling

- Based on data as much as possible
- W/Z+jets production
  - Estimated from MC/data
    - Distributions from MC
    - Normalization from pre-tagged sample
    - Flavor fractions from NLO



- Multijet events (misidentified lepton)
  - Estimated from data
- Top pair production
  - Estimated from MC
- Diboson (WZ, WW)
  - Estimated from MC



Event Yield in 370 pb^{-1}

	s-channel	t-channel
Cut acceptance	23%	22%
b-tag efficiency	54%	38%
Signal yield	9.5	15.0
Bkgnd yield	452	
Data	443	
Signal/bkgnd	1:50	1:30



Pre-tagged
10,000 events

=1 b-tag
367 events

=2 b-tags
76 events







- Wjj
- Wbb
- $t\bar{t}$
- WW/WZ
- t-channel
- s-channel

Separating Signal from Backgrounds

- Four analysis methods used so far
  - Each using the same analysis strategy

Cut-Based



Decision Trees



Neural Networks



Likelihood



Analysis Strategy



Analysis Strategy



Analysis Strategy



Discriminating Variables

- Object kinematics
  - Jet p_T for different jets
    - Tagged, untagged,...
- Event kinematics
  - H (total energy)
  - H_T (transverse energy)
  - M (invariant mass)
  - M_T (transverse mass)
  - Summing over various objects in the event
- Angular variables
  - Jet-jet separation
  - Jet pseudorapidity (t-channel)
  - Top quark spin
  - Sphericity, aplanarity



25 variables total



Neural Networks

Input Nodes: One for each variable x_i



Output Node: linear combination of hidden nodes

$$f(\vec{x}) = \Sigma\, w'_k\, n_k(\vec{x},\vec{w}_k)$$



Hidden Nodes: Each is a sigmoid dependent on the input variables

$$n_k(\vec{x},\vec{w}_k) = \frac{1}{1 + e^{-\Sigma\, w_{ik}\, x_i}}$$



Neural Network Output



Neural Network Result

- Dominant systematics
  - Jet energy scale, b-ID: 10%
  - Background norm: 15-20%
  - Object ID: 5%
- Total systematic

	=1 tag	≥2 tags
Signal acceptance	15%	25%
Background sum	10%	26%



Expected/Observed limit:

σ_s < 4.5 / 6.4 pb

σ_t < 5.8 / 5.0 pb

Published in PLB 622, 265 (2005)

Sensitivity to non-SM Single Top



using only muon channel data

using only electron channel data

V_{tb} Coupling Limit

- Both the s-channel and t-channel cross sections are proportional to $|V_{tb}|^2$
  - Combine neural networks in a 2d likelihood function



  - Need to assume SM ratio of cross sections

Bayesian Posterior density for $|V_{tb}|^2$



2. Likelihood Analysis (370 pb^{-1})



- Using jet lifetime tagger
  - Asymmetric tag requirements
    - At least one tight tag
    - Loose tagger cut for double tags



Likelihood Result

- Systematic uncertainties
  - b-tag modeling: 6-17%
  - Jet energy scale: 5%
  - Trigger modeling: 5%
  - Theory tt XS: 18%
  -



Expected/Observed limit:

σ_s < 3.3 / 5.0 pb

σ_t < 4.3 / 4.4 pb



Summary

	s-channel	t-channel
NLO cross section	0.88 pb	1.98 pb
	95% CL upper cross section limits [pb]	
DØ Run I	17	22
CDF Run II ($160pb^{-1}$)	13.6	10.1
Neural Network ($230pb^{-1}$)	6.4	5.0
Likelihood ($370pb^{-1}$)	5.0	4.4

Outlook

The future is here

Tevatron Single Top Future



Improvements

- b-tagging
  - More signal
- Jet energy resolution
  - Better top mass reconstruction

Prospects

- Observe single top production in Run II
  - Discover new physics (if it's there)
- Measure $|V_{tb}|$ to ~10%

Tevatron luminosity projection



Tevatron luminosity projection



Outlook

Top is the gateway quark

Energy Frontier of the Future: LHC





SM Single Top Production at the LHC

s-channel



t-channel



associated production



- Observe three single top production modes separately
- Measure V_{tb} to few %
- Study spin correlations

LHC as a Higgs Discovery Machine

Neutral Higgs

Dominant Higgs production mechanism at the LHC:



Top-Higgs Yukawa coupling



Discover with ~30fb^{-1}

Supersymmetry: Charged Higgs

Heavy H^+:



Discovery possible with ~30fb^{-1}

Light H^+:



Discover with ~30fb^{-1}

Top quark Yukawa coupling

Precision coupling measurement at a future e^+e^- collider



International Linear Collider

Conclusions

- Single top quark production is a key observation for the Tevatron
- DØ Run II single top analysis going well
  - World's best limits: $\sigma_s < 5.0$ pb, $\sigma_t < 4.4$ pb
  - Already reaching sensitivity to new physics
- Single Top is about to get really exciting in Run II
  - Will be close to discovery in ~ 1 year
- Top quark couplings will be studied in detail at future colliders