

Search for flavor changing neutral currents in decays of top quarks

V.M. Abazov,³⁵ B. Abbott,⁷³ B.S. Acharya,²⁹ M. Adams,⁴⁹ T. Adams,⁴⁷ G.D. Alexeev,³⁵ G. Alkhazov,³⁹ A. Alton^a,⁶¹ G. Alverson,⁶⁰ G.A. Alves,² L.S. Ancu,³⁴ M. Aoki,⁴⁸ M. Arov,⁵⁸ A. Askew,⁴⁷ B. Åsman,⁴¹ O. Atramentov,⁶⁵ C. Avila,⁸ J. BackusMayes,⁸⁰ F. Badaud,¹³ L. Bagby,⁴⁸ B. Baldin,⁴⁸ D.V. Bandurin,⁴⁷ S. Banerjee,²⁹ E. Barberis,⁶⁰ P. Baringer,⁵⁶ J. Barreto,³ J.F. Bartlett,⁴⁸ U. Bassler,¹⁸ V. Bazterra,⁴⁹ S. Beale,⁶ A. Bean,⁵⁶ M. Begalli,³ M. Begel,⁷¹ C. Belanger-Champagne,⁴¹ L. Bellantoni,⁴⁸ S.B. Beri,²⁷ G. Bernardi,¹⁷ R. Bernhard,²² I. Bertram,⁴² M. Besançon,¹⁸ R. Beuselinck,⁴³ V.A. Bezzubov,³⁸ P.C. Bhat,⁴⁸ V. Bhatnagar,²⁷ G. Blazey,⁵⁰ S. Blessing,⁴⁷ K. Bloom,⁶⁴ A. Boehnlein,⁴⁸ D. Boline,⁷⁰ T.A. Bolton,⁵⁷ E.E. Boos,³⁷ G. Borissov,⁴² T. Bose,⁵⁹ A. Brandt,⁷⁶ O. Brandt,²³ R. Brock,⁶² G. Brooijmans,⁶⁸ A. Bross,⁴⁸ D. Brown,¹⁷ J. Brown,¹⁷ X.B. Bu,⁴⁸ M. Buehler,⁷⁹ V. Buescher,²⁴ V. Bunichev,³⁷ S. Burdin^b,⁴² T.H. Burnett,⁸⁰ C.P. Buszello,⁴¹ B. Calpas,¹⁵ E. Camacho-Pérez,³² M.A. Carrasco-Lizarraga,⁵⁶ B.C.K. Casey,⁴⁸ H. Castilla-Valdez,³² S. Chakrabarti,⁷⁰ D. Chakraborty,⁵⁰ K.M. Chan,⁵⁴ A. Chandra,⁷⁸ G. Chen,⁵⁶ S. Chevalier-Théry,¹⁸ D.K. Cho,⁷⁵ S.W. Cho,³¹ S. Choi,³¹ B. Choudhary,²⁸ T. Christoudias,⁴³ S. Cihangir,⁴⁸ D. Claes,⁶⁴ J. Clutter,⁵⁶ M. Cooke,⁴⁸ W.E. Cooper,⁴⁸ M. Corcoran,⁷⁸ F. Couderc,¹⁸ M.-C. Cousinou,¹⁵ A. Croc,¹⁸ D. Cutts,⁷⁵ A. Das,⁴⁵ G. Davies,⁴³ K. De,⁷⁶ S.J. de Jong,³⁴ E. De La Cruz-Burelo,³² F. Déliot,¹⁸ M. Demarteau,⁴⁸ R. Demina,⁶⁹ D. Denisov,⁴⁸ S.P. Denisov,³⁸ S. Desai,⁴⁸ K. DeVaughan,⁶⁴ H.T. Diehl,⁴⁸ M. Diesburg,⁴⁸ A. Dominguez,⁶⁴ T. Dorland,⁸⁰ A. Dubey,²⁸ L.V. Dudko,³⁷ D. Duggan,⁶⁵ A. Duperrin,¹⁵ S. Dutt,²⁷ A. Dyshkant,⁵⁰ M. Eads,⁶⁴ D. Edmunds,⁶² J. Ellison,⁴⁶ V.D. Elvira,⁴⁸ Y. Enari,¹⁷ H. Evans,⁵² A. Evdokimov,⁷¹ V.N. Evdokimov,³⁸ G. Facini,⁶⁰ T. Ferbel,⁶⁹ F. Fiedler,²⁴ F. Filthaut,³⁴ W. Fisher,⁶² H.E. Fisk,⁴⁸ M. Fortner,⁵⁰ H. Fox,⁴² S. Fuess,⁴⁸ T. Gadfort,⁷¹ A. Garcia-Bellido,⁶⁹ V. Gavrilov,³⁶ P. Gay,¹³ W. Geist,¹⁹ W. Geng,^{15,62} D. Gerbaudo,⁶⁶ C.E. Gerber,⁴⁹ Y. Gershtein,⁶⁵ G. Ginther,^{48,69} G. Golovanov,³⁵ A. Goussiou,⁸⁰ P.D. Grannis,⁷⁰ S. Greder,¹⁹ H. Greenlee,⁴⁸ Z.D. Greenwood,⁵⁸ E.M. Gregores,⁴ G. Grenier,²⁰ Ph. Gris,¹³ J.-F. Grivaz,¹⁶ A. Grohsjean,¹⁸ S. Grünendahl,⁴⁸ M.W. Grünewald,³⁰ T. Guillemin,¹⁶ F. Guo,⁷⁰ G. Gutierrez,⁴⁸ P. Gutierrez,⁷³ A. Haas^c,⁶⁸ S. Hagopian,⁴⁷ J. Haley,⁶⁰ L. Han,⁷ K. Harder,⁴⁴ A. Harel,⁶⁹ J.M. Hauptman,⁵⁵ J. Hays,⁴³ T. Head,⁴⁴ T. Hebbeker,²¹ D. Hedin,⁵⁰ H. Hegab,⁷⁴ A.P. Heinson,⁴⁶ U. Heintz,⁷⁵ C. Hensel,²³ I. Heredia-De La Cruz,³² K. Herner,⁶¹ G. Hesketh^d,⁴⁴ M.D. Hildreth,⁵⁴ R. Hirosky,⁷⁹ T. Hoang,⁴⁷ J.D. Hobbs,⁷⁰ B. Hoeneisen,¹² M. Hohlfield,²⁴ Z. Hubacek,^{10,18} N. Huske,¹⁷ V. Hynek,¹⁰ I. Iashvili,⁶⁷ R. Illingworth,⁴⁸ A.S. Ito,⁴⁸ S. Jabeen,⁷⁵ M. Jaffré,¹⁶ D. Jamin,¹⁵ A. Jayasinghe,⁷³ R. Jesik,⁴³ K. Johns,⁴⁵ M. Johnson,⁴⁸ D. Johnston,⁶⁴ A. Jonckheere,⁴⁸ P. Jonsson,⁴³ J. Joshi,²⁷ A. Juste,⁴⁰ K. Kaadze,⁵⁷ E. Kajfasz,¹⁵ D. Karmanov,³⁷ P.A. Kasper,⁴⁸ I. Katsanos,⁶⁴ R. Kehoe,⁷⁷ S. Kermiche,¹⁵ N. Khalatyan,⁴⁸ A. Khanov,⁷⁴ A. Kharchilava,⁶⁷ Y.N. Kharzheev,³⁵ D. Khatidze,⁷⁵ M.H. Kirby,⁵¹ J.M. Kohli,²⁷ A.V. Kozelov,³⁸ J. Kraus,⁶² S. Kulikov,³⁸ A. Kumar,⁶⁷ A. Kupco,¹¹ T. Kurča,²⁰ V.A. Kuzmin,³⁷ J. Kvita,⁹ S. Lammers,⁵² G. Landsberg,⁷⁵ P. Lebrun,²⁰ H.S. Lee,³¹ S.W. Lee,⁵⁵ W.M. Lee,⁴⁸ J. Lellouch,¹⁷ L. Li,⁴⁶ Q.Z. Li,⁴⁸ S.M. Lietti,⁵ J.K. Lim,³¹ D. Lincoln,⁴⁸ J. Linnemann,⁶² V.V. Lipaev,³⁸ R. Lipton,⁴⁸ Y. Liu,⁷ Z. Liu,⁶ A. Lobodenko,³⁹ M. Lokajicek,¹¹ R. Lopes de Sa,⁷⁰ H.J. Lubatti,⁸⁰ R. Luna-Garcia^e,³² A.L. Lyon,⁴⁸ A.K.A. Maciel,² D. Mackin,⁷⁸ R. Madar,¹⁸ R. Magaña-Villalba,³² S. Malik,⁶⁴ V.L. Malyshev,³⁵ Y. Maravin,⁵⁷ J. Martínez-Ortega,³² R. McCarthy,⁷⁰ C.L. McGivern,⁵⁶ M.M. Meijer,³⁴ A. Melnitchouk,⁶³ D. Menezes,⁵⁰ P.G. Mercadante,⁴ M. Merkin,³⁷ A. Meyer,²¹ J. Meyer,²³ F. Miconi,¹⁹ N.K. Mondal,²⁹ G.S. Muanza,¹⁵ M. Mulhearn,⁷⁹ E. Nagy,¹⁵ M. Naimuddin,²⁸ M. Narain,⁷⁵ R. Nayyar,²⁸ H.A. Neal,⁶¹ J.P. Negret,⁸ P. Neustroev,³⁹ S.F. Novaes,⁵ T. Nunnemann,²⁵ G. Obrant,³⁹ J. Orduna,⁷⁸ N. Osman,¹⁵ J. Osta,⁵⁴ G.J. Otero y Garzón,¹ M. Padilla,⁴⁶ A. Pal,⁷⁶ M. Pangilinan,⁷⁵ N. Parashar,⁵³ V. Parihar,⁷⁵ S.K. Park,³¹ J. Parsons,⁶⁸ R. Partridge^c,⁷⁵ N. Parua,⁵² A. Patwa,⁷¹ B. Penning,⁴⁸ M. Perfilov,³⁷ K. Peters,⁴⁴ Y. Peters,⁴⁴ K. Petridis,⁴⁴ G. Petrillo,⁶⁹ P. Pétrouff,¹⁶ R. Piegai,¹ J. Piper,⁶² M.-A. Pleier,⁷¹ P.L.M. Podesta-Lerma^f,³² V.M. Podstavkov,⁴⁸ M.-E. Pol,² P. Polozov,³⁶ A.V. Popov,³⁸ M. Prewitt,⁷⁸ D. Price,⁵² N. Prokopenko,³⁸ S. Protopopescu,⁷¹ J. Qian,⁶¹ A. Quadt,²³ B. Quinn,⁶³ M.S. Rangel,² K. Ranjan,²⁸ P.N. Ratoff,⁴² I. Razumov,³⁸ P. Renkel,⁷⁷ M. Rijssenbeek,⁷⁰ I. Ripp-Baudot,¹⁹ F. Rizatdinova,⁷⁴ M. Rominsky,⁴⁸ A. Ross,⁴² C. Royon,¹⁸ P. Rubinov,⁴⁸ R. Ruchti,⁵⁴ G. Safronov,³⁶ G. Sajot,¹⁴ P. Salcido,⁵⁰ A. Sánchez-Hernández,³² M.P. Sanders,²⁵ B. Sanghi,⁴⁸ A.S. Santos,⁵ G. Savage,⁴⁸ L. Sawyer,⁵⁸ T. Scanlon,⁴³ R.D. Schamberger,⁷⁰ Y. Scheglov,³⁹ H. Schellman,⁵¹ T. Schliephake,²⁶ S. Schlobohm,⁸⁰ C. Schwanenberger,⁴⁴ R. Schwenhorst,⁶² J. Sekaric,⁵⁶ H. Severini,⁷³ E. Shabalina,²³ V. Shary,¹⁸ A.A. Shchukin,³⁸ R.K. Shivpuri,²⁸ V. Simak,¹⁰ V. Sirotenko,⁴⁸ P. Skubic,⁷³ P. Slatery,⁶⁹ D. Smirnov,⁵⁴ K.J. Smith,⁶⁷ G.R. Snow,⁶⁴ J. Snow,⁷² S. Snyder,⁷¹ S. Söldner-Rembold,⁴⁴ L. Sonnenschein,²¹ K. Soustruznik,⁹ J. Stark,¹⁴ V. Stolin,³⁶ D.A. Stoyanova,³⁸ M. Strauss,⁷³

D. Strom,⁴⁹ L. Stutte,⁴⁸ L. Suter,⁴⁴ P. Svoisky,⁷³ M. Takahashi,⁴⁴ A. Tanasijczuk,¹ W. Taylor,⁶ M. Titov,¹⁸ V.V. Tokmenin,³⁵ Y.-T. Tsai,⁶⁹ D. Tsybychev,⁷⁰ B. Tuchming,¹⁸ C. Tully,⁶⁶ P.M. Tuts,⁶⁸ L. Uvarov,³⁹ S. Uvarov,³⁹ S. Uzunyan,⁵⁰ R. Van Kooten,⁵² W.M. van Leeuwen,³³ N. Varelas,⁴⁹ E.W. Varnes,⁴⁵ I.A. Vasilyev,³⁸ P. Verdier,²⁰ L.S. Vertogradov,³⁵ M. Verzocchi,⁴⁸ M. Vesterinen,⁴⁴ D. Vilanova,¹⁸ P. Vint,⁴³ P. Vokac,¹⁰ H.D. Wahl,⁴⁷ M.H.L.S. Wang,⁶⁹ J. Warchol,⁵⁴ G. Watts,⁸⁰ M. Wayne,⁵⁴ M. Weber,^{9,48} L. Welty-Rieger,⁵¹ A. White,⁷⁶ D. Wicke,²⁶ M.R.J. Williams,⁴² G.W. Wilson,⁵⁶ M. Wobisch,⁵⁸ D.R. Wood,⁶⁰ T.R. Wyatt,⁴⁴ Y. Xie,⁴⁸ C. Xu,⁶¹ S. Yacoob,⁵¹ R. Yamada,⁴⁸ W.-C. Yang,⁴⁴ T. Yasuda,⁴⁸ Y.A. Yatsunenko,³⁵ Z. Ye,⁴⁸ H. Yin,⁴⁸ K. Yip,⁷¹ S.W. Youn,⁴⁸ J. Yu,⁷⁶ S. Zelitch,⁷⁹ T. Zhao,⁸⁰ B. Zhou,⁶¹ J. Zhu,⁶¹ M. Zielinski,⁶⁹ D. Zieminska,⁵² and L. Zivkovic⁷⁵

(The D0 Collaboration*)

¹Universidad de Buenos Aires, Buenos Aires, Argentina

²LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

³Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

⁴Universidade Federal do ABC, Santo André, Brazil

⁵Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil

⁶Simon Fraser University, Vancouver, British Columbia, and York University, Toronto, Ontario, Canada

⁷University of Science and Technology of China, Hefei, People's Republic of China

⁸Universidad de los Andes, Bogotá, Colombia

⁹Charles University, Faculty of Mathematics and Physics,
Center for Particle Physics, Prague, Czech Republic

¹⁰Czech Technical University in Prague, Prague, Czech Republic

¹¹Center for Particle Physics, Institute of Physics,
Academy of Sciences of the Czech Republic, Prague, Czech Republic

¹²Universidad San Francisco de Quito, Quito, Ecuador

¹³LPC, Université Blaise Pascal, CNRS/IN2P3, Clermont, France

¹⁴LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3,
Institut National Polytechnique de Grenoble, Grenoble, France

¹⁵CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France

¹⁶LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France

¹⁷LPNHE, Universités Paris VI and VII, CNRS/IN2P3, Paris, France

¹⁸CEA, Irfu, SPP, Saclay, France

¹⁹IPHC, Université de Strasbourg, CNRS/IN2P3, Strasbourg, France

²⁰IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France and Université de Lyon, Lyon, France

²¹III. Physikalisches Institut A, RWTH Aachen University, Aachen, Germany

²²Physikalisches Institut, Universität Freiburg, Freiburg, Germany

²³II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen, Germany

²⁴Institut für Physik, Universität Mainz, Mainz, Germany

²⁵Ludwig-Maximilians-Universität München, München, Germany

²⁶Fachbereich Physik, Bergische Universität Wuppertal, Wuppertal, Germany

²⁷Panjab University, Chandigarh, India

²⁸Delhi University, Delhi, India

²⁹Tata Institute of Fundamental Research, Mumbai, India

³⁰University College Dublin, Dublin, Ireland

³¹Korea Detector Laboratory, Korea University, Seoul, Korea

³²CINVESTAV, Mexico City, Mexico

³³FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands

³⁴Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands

³⁵Joint Institute for Nuclear Research, Dubna, Russia

³⁶Institute for Theoretical and Experimental Physics, Moscow, Russia

³⁷Moscow State University, Moscow, Russia

³⁸Institute for High Energy Physics, Protvino, Russia

³⁹Petersburg Nuclear Physics Institute, St. Petersburg, Russia

⁴⁰Institució Catalana de Recerca i Estudis Avançats (ICREA) and Institut de Física d'Altes Energies (IFAE), Barcelona, Spain

⁴¹Stockholm University, Stockholm and Uppsala University, Uppsala, Sweden

⁴²Lancaster University, Lancaster LA1 4YB, United Kingdom

⁴³Imperial College London, London SW7 2AZ, United Kingdom

⁴⁴The University of Manchester, Manchester M13 9PL, United Kingdom

⁴⁵University of Arizona, Tucson, Arizona 85721, USA

⁴⁶University of California Riverside, Riverside, California 92521, USA

⁴⁷Florida State University, Tallahassee, Florida 32306, USA

⁴⁸Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA

⁴⁹University of Illinois at Chicago, Chicago, Illinois 60607, USA

⁵⁰Northern Illinois University, DeKalb, Illinois 60115, USA

- ⁵¹Northwestern University, Evanston, Illinois 60208, USA
⁵²Indiana University, Bloomington, Indiana 47405, USA
⁵³Purdue University Calumet, Hammond, Indiana 46323, USA
⁵⁴University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵⁵Iowa State University, Ames, Iowa 50011, USA
⁵⁶University of Kansas, Lawrence, Kansas 66045, USA
⁵⁷Kansas State University, Manhattan, Kansas 66506, USA
⁵⁸Louisiana Tech University, Ruston, Louisiana 71272, USA
⁵⁹Boston University, Boston, Massachusetts 02215, USA
⁶⁰Northeastern University, Boston, Massachusetts 02115, USA
⁶¹University of Michigan, Ann Arbor, Michigan 48109, USA
⁶²Michigan State University, East Lansing, Michigan 48824, USA
⁶³University of Mississippi, University, Mississippi 38677, USA
⁶⁴University of Nebraska, Lincoln, Nebraska 68588, USA
⁶⁵Rutgers University, Piscataway, New Jersey 08855, USA
⁶⁶Princeton University, Princeton, New Jersey 08544, USA
⁶⁷State University of New York, Buffalo, New York 14260, USA
⁶⁸Columbia University, New York, New York 10027, USA
⁶⁹University of Rochester, Rochester, New York 14627, USA
⁷⁰State University of New York, Stony Brook, New York 11794, USA
⁷¹Brookhaven National Laboratory, Upton, New York 11973, USA
⁷²Langston University, Langston, Oklahoma 73050, USA
⁷³University of Oklahoma, Norman, Oklahoma 73019, USA
⁷⁴Oklahoma State University, Stillwater, Oklahoma 74078, USA
⁷⁵Brown University, Providence, Rhode Island 02912, USA
⁷⁶University of Texas, Arlington, Texas 76019, USA
⁷⁷Southern Methodist University, Dallas, Texas 75275, USA
⁷⁸Rice University, Houston, Texas 77005, USA
⁷⁹University of Virginia, Charlottesville, Virginia 22901, USA
⁸⁰University of Washington, Seattle, Washington 98195, USA
- (Dated: March 23, 2011)

We present a search for flavor changing neutral currents in decays of top quarks. The analysis is based on a search for $t\bar{t} \rightarrow \ell'\nu\ell\ell + \text{jets}$ ($\ell, \ell' = e, \mu$) final states using 4.1 fb^{-1} of integrated luminosity of $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$. We extract limits on the branching ratio $B(t \rightarrow Zq)$ ($q = u, c$ quarks), assuming anomalous tuZ or tcZ couplings. We do not observe any sign of such anomalous coupling and set a limit of $B < 3.2\%$ at 95% C.L.

PACS numbers: 12.60.Cn, 13.85.Qk, 14.65.Ha, 12.15.Mm

Flavor changing neutral currents (FCNC) allow for transitions between quarks of different flavor but same electric charge. FCNC are sensitive indicators of physics beyond the standard model (BSM), because they are suppressed in the standard model (SM).

In this paper, we search for FCNC decays of the top (t) quark [1]. Within the SM the top quark decays into a W boson and a b quark with a rate proportional to the Cabibbo-Kobayashi-Maskawa (CKM) matrix element squared, $|V_{tb}|^2$ [2]. Under the assumption of three fermion families and a unitary 3×3 CKM matrix, the $|V_{tb}|$ element is severely constrained to $|V_{tb}| = 0.999152^{+0.000030}_{-0.000045}$ [3]. While the SM branching frac-

tion for $t \rightarrow Zq$ ($q = u, c$ quarks) is predicted to be $\approx 10^{-14}$ [4], supersymmetric extensions of the SM with or without R -parity violation, or quark compositeness predict branching fractions as high as $\approx 10^{-4}$ [4-6]. The observation of the FCNC decay $t \rightarrow Zq$ would therefore provide evidence of contributions from BSM physics.

We analyze top-pair production ($t\bar{t}$), where either one or both of the top quarks decay via $t \rightarrow Zq$ or their charge conjugates (hereafter implied). Any top quark that does not decay via $t \rightarrow Zq$ is assumed to decay via $t \rightarrow Wb$. We assume that the $t \rightarrow Zq$ decay is generated by an anomalous FCNC term added to the SM Lagrangian

$$\mathcal{L}_{\text{FCNC}} = \frac{e}{2 \sin \theta_W \cos \theta_W} \bar{t} \gamma_\mu (v_{tqZ} - a_{tqZ} \gamma_5) q Z^\mu + h.c., \quad (1)$$

where q , t , and Z are the quantum fields for up or charm quarks, top quarks, and for the Z boson, respectively, e is the electric charge and θ_W the Weinberg angle. We thereby introduce dimension-4 vector, v_{tqZ} , and axial vector, a_{tqZ} , couplings as defined in [7].

*with visitors from ^aAugustana College, Sioux Falls, SD, USA, ^bThe University of Liverpool, Liverpool, UK, ^cSLAC, Menlo Park, CA, USA, ^dUniversity College London, London, UK, ^eCentro de Investigacion en Computacion - IPN, Mexico City, Mexico, ^fEFCM, Universidad Autonoma de Sinaloa, Culiacán, Mexico, and ^gUniversität Bern, Bern, Switzerland.




Figure 1: Lowest-order diagram for FCNC $t\bar{t} \rightarrow WbZq'$ production, where q' can be either a u or c quark, and the W and Z bosons decay leptonically.

We investigate channels where the W and Z bosons decay leptonically, as shown in Fig. 1. The u , c , and b quarks subsequently hadronize, giving rise to a final state with three charged leptons ($\ell = e, \mu$), an imbalance in momentum transverse to the $p\bar{p}$ collision axis (\cancel{E}_T , from the escaping neutrino in the $W \rightarrow \ell\nu$ decay), and jets.

This is the first search for FCNC in $t\bar{t}$ decays with trilepton final states. This mode provides a distinct signature with low background, albeit at the cost of statistical power. Previous studies have been done to search for FCNC processes in meson decays, i.e., $b \rightarrow Zs$ in $B^+ \rightarrow K^{*+}\ell^+\ell^-$ [8–10] or $s \rightarrow Zd$ in $K^+ \rightarrow \pi^+\nu\bar{\nu}$ [11]. Using the $D^+ \rightarrow \pi^+\mu^+\mu^-$ final state in 1.3 fb^{-1} of integrated luminosity, the D0 Collaboration has set the best branching ratio (B) limits on the FCNC $c \rightarrow Zu$ process at $B(D^+ \rightarrow \pi^+\mu^+\mu^-) < 3.9 \times 10^{-6}$ at 90% C.L. [12]. There are theoretical arguments as to why top quark decays may be the best way to study flavor violating couplings of mass-dependent interactions [13, 14]. FCNC tqZ and $tq\gamma$ couplings have been studied by the CERN e^+e^- Collider (LEP), DESY ep Collider (HERA), and Tevatron collider experiments [15–19]. The D0 Collaboration has recently published limits on the branching ratios determined from FCNC gluon-quark couplings using single top quark final states [20]. The 95% C.L. upper limit on the branching ratio of $t \rightarrow Zq$ from the CDF Collaboration uses 1.9 fb^{-1} of integrated luminosity, assumes a top quark mass of $m_t = 175 \text{ GeV}$ and uses the measured cross section of $\sigma_{t\bar{t}} = 8.8 \pm 1.1 \text{ pb}$ [19]. This result excludes branching ratios of $B(t \rightarrow Zq) > 3.7\%$, with an expected limit of $5.0\% \pm 2.2\%$. To obtain these results, CDF exploited the two lepton plus four jet final state. This signature occurs when one of the pair-produced top quarks decays via FCNC to Zq , followed by the decay $Z \rightarrow ee$ or $Z \rightarrow \mu\mu$. The other top quark decays to Wb , followed by the hadronic decay of the W boson. This dilepton signature suffers from large background, but profits from more statistics relative to the trilepton final states investigated in this Letter.

This analysis is based on the measurement of the WZ production cross section in $\ell\nu\ell\ell$ final states [21] us-

ing 4.1 fb^{-1} of integrated luminosity of $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$. We require the presence of one or more jets in the final state and investigate observables that are sensitive to the signal topology in order to select events with $WZ \rightarrow \ell\nu\ell\ell$ decays that originate from the pair production of top quarks.

A detailed description of the D0 detector can be found elsewhere [22]. Here, we give a brief overview of the main sub-systems of the detector. The innermost part is a central tracking system surrounded by a 1.9 T superconducting solenoidal magnet. The two components of the central tracking system, a silicon microstrip tracker and a central fiber tracker, are used to reconstruct $p\bar{p}$ interaction vertices and provide the measurement of the momentum of charged particles within the pseudorapidity range of $|\eta| < 2$ (with η defined relative to the center of the detector). The tracking system and magnet are followed by the calorimetry that consists of central (CC) and end (EC) electromagnetic and hadronic uranium-liquid argon sampling calorimeters, and an intercryostat detector (ICD). The CC and two EC calorimeters cover $|\eta| < 1.1$ and $1.5 < |\eta| < 4.2$, respectively, while the ICD covers for $1.1 < |\eta| < 1.4$. The calorimeter measures energies of hadrons, electrons, and photons. The muon system consists of a layer of drift tubes and scintillation counters inside a 1.8 T toroidal magnet. Two similar layers are outside of the toroidal magnet and the entire system covers $|\eta| < 2$.

An electron is identified from the properties of clusters of energy deposited in the CC, EC, or ICD that match a track reconstructed in the central tracker. Because of the lack of far forward coverage of the tracker, we define EC electrons only within $1.5 < |\eta| < 2.5$. The calorimeter clusters in the CC and EC are required to pass the isolation cut

$$\frac{E_{\text{tot}}(\Delta\mathcal{R} < 0.4) - E_{\text{EM}}(\Delta\mathcal{R} < 0.2)}{E_{\text{EM}}(\Delta\mathcal{R} < 0.2)} < 0.1$$

for the "loose" electrons and < 0.07 for "tight" electrons, where E_{tot} is the total energy in the EM and hadronic calorimeters, E_{EM} is the energy found in the EM calorimeter only, and $\Delta\mathcal{R} = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2}$. For the intercryostat region (ICR), $1.1 < |\eta| < 1.5$, we form clusters from the energy deposits in the CC, ICD, or EC detectors. These clusters are identified as electrons if they pass a neural network requirement that is based on the characteristics of the shower and associated track information. A muon candidate is reconstructed from track segments within the muon system that are matched to a track reconstructed in the central tracker. The trajectory of the muon candidate must be isolated from other tracks in the tracker and the energy deposition in the calorimeter. Jets are reconstructed from the energy deposited in the CC and EC calorimeters, using the "Run II midpoint cone" algorithm [23] of size $\Delta\mathcal{R} = 0.5$, within $|\eta| < 2.5$, where ϕ is the azimuthal angle.

Monte Carlo (MC) samples of WZ and ZZ back-

ground events are produced using the PYTHIA generator [24]. The production of the W and Z bosons in association with jets (W +jets, Z +jets), collectively referred to as V +jets, as well as $t\bar{t}$ processes are generated using ALPGEN [25] interfaced with PYTHIA for parton evolution and hadronization. In all samples the CTEQ6L1 parton distribution function (PDF) set is used, along with $m_t = 172.5$ GeV. The $t\bar{t}$ cross section is set to the SM value at this top quark mass, i.e., $\sigma_{t\bar{t}} = 7.46_{-0.67}^{+0.48}$ pb [26]. This uncertainty is mainly due to the scale dependence, PDFs, and the experimental uncertainty on m_t [27].

All MC samples are passed through a GEANT [28] simulation of the D0 detector and overlaid with data events from random beam crossings to account for the underlying event. The samples are then corrected for the luminosity dependence of the trigger, reconstruction efficiencies in data, and the beam position. All MC samples are normalized to the luminosity in data using next-to-leading order (NLO) calculations of the cross sections, and are subject to the same selection criteria as applied to data.

The signal process is generated using the PYTHIA generator with the decay $t \rightarrow Zq$ added. The Z boson helicity is implemented by reweighting an angular distribution of the positively charged lepton in the decay $t \rightarrow Zq \rightarrow \ell^+ \ell^- q$ using CompHEP [29], modified by the addition of the Lagrangian of Eq. 1. The variable $\cos\theta^*$ used for the reweighting is defined by the angle θ^* between the Z boson's momentum in the top quark rest frame and the momentum of the positively charged lepton in the Z boson rest frame. We assume in the analysis that the vector and axial vector couplings, as introduced in Eq. 1, are identical to the corresponding couplings for neutral currents (NC) in the SM, i.e., $v_{tuZ} = 1/2 - 4/3 \sin^2 \theta_W = 0.192$ and $a_{tuZ} = 1/2$, where $\sin^2 \theta_W = 0.231$. To study the influence of different values of the couplings, we also analyse the following cases: (i) $v_{tuZ} = 1, a_{tuZ} = 0$; (ii) $v_{tuZ} = 0, a_{tuZ} = 1$; and (iii) $v_{tuZ} = a_{tuZ} = 1/\sqrt{2}$. The first two give identical results. The difference obtained by using cases (i/ii), (iii), and using the values of the SM NC couplings is included as systematic uncertainty. Therefore, our result is independent of the actual values of v_{tqZ} and a_{tqZ} . Since we do not distinguish c and u quark jets our results are valid also for u and c quarks separately.

The total selection efficiency, calculated as a function of $B = \Gamma(t \rightarrow Zq)/(\Gamma(t \rightarrow Wb) + \Gamma(t \rightarrow Zq))$, can be written as

$$\begin{aligned} \epsilon_{t\bar{t}} = & (1 - B)^2 \cdot \epsilon_{t\bar{t} \rightarrow W^+ b W^- \bar{b}} \\ & + 2B(1 - B) \cdot \epsilon_{t\bar{t} \rightarrow Zq W^- \bar{b}} + B^2 \cdot \epsilon_{t\bar{t} \rightarrow Zq Zq}, \end{aligned} \quad (2)$$

where the efficiency $\epsilon_{t\bar{t} \rightarrow W^+ b W^- \bar{b}}$ for the SM $t\bar{t}$ background contribution is used, along with the efficiencies $\epsilon_{t\bar{t} \rightarrow Zq W^- \bar{b}}$ and $\epsilon_{t\bar{t} \rightarrow Zq Zq}$ that include the FCNC top quark decays.

We consider four independent decay signatures: $eee + \cancel{E}_T + X$, $ee\mu + \cancel{E}_T + X$, $\mu\mu e + \cancel{E}_T + X$, and $\mu\mu\mu + \cancel{E}_T + X$,

where X is any number of jets. We require the events to have at least three lepton candidates with transverse momentum $p_T > 15$ GeV that originate from the same $p\bar{p}$ interaction vertex and are separated from each other by $\Delta\mathcal{R} > 0.5$. The event must also have $\cancel{E}_T > 20$ GeV, which is calculated from the energy found in the calorimeter cells and p_T corrected for any muons reconstructed in the event. Furthermore, all energy corrections applied to electrons and jets are propagated through to the \cancel{E}_T .

Events are selected using triggers based on electrons and muons. There are several high- p_T leptons from the decay of the heavy gauge bosons providing a total trigger efficiency for all signatures of $98\% \pm 2\%$.

To identify the leptons from the Z boson decay, we consider only pairs of electrons or muons, additionally requiring them to have opposite electric charges. If no lepton pair is found within the invariant mass intervals of 74–108 GeV (ee), 65–115 GeV ($\mu\mu$) or 60–120 GeV (ee , with one electron in the ICR) the event is rejected, else, the pair that has an invariant mass closest to the Z boson mass M_Z is selected as the Z boson. The lepton with the highest p_T of the remaining muons or CC/EC electrons in the event is selected as originating from the W boson decay. From simulation, this assignment of the three charged leptons to Z and W bosons is found to be $\approx 100\%$ correct for $ee\mu$ and $\mu\mu e$, and about 92% and 89% for the eee and $\mu\mu\mu$ channels, respectively.

If jets are present, they are required to have $p_T > 20$ GeV and $|\eta| < 2.5$. We also require that jets are separated from electrons by $\Delta\mathcal{R} > 0.5$. There is no fixed separation cut between the muon and jets but the muon isolation requirement rejects most muons within $\Delta\mathcal{R} < 0.4$ of a jet. Thresholds in the selection criteria are the same as in Ref. [21] and the acceptance multiplied by efficiency results are summarized in Table I for the FCNC signal.

In addition to SM WZ production, the other major background is from processes with a Z boson and an additional object misidentified as the lepton from the W boson decay (e.g., from Z +jets, ZZ , and $Z\gamma$). A small background contribution is expected from processes such as W +jets and SM $t\bar{t}$ production. The WZ , ZZ , and $t\bar{t}$ backgrounds are estimated from the simulation, while the V +jets and $Z\gamma$ backgrounds are estimated using data-driven methods.

One or more jets in V +jets events can be misidentified as a lepton from W or Z boson decays. To estimate this contribution, we define a *false* lepton category for electrons and muons. A *false* electron is required to have most of its energy deposited in the electromagnetic part of the calorimeter and satisfy calorimeter isolation criteria for electrons, but have a shower shape inconsistent with that of an electron. A muon candidate is categorized as *false* if it fails isolation criteria, as determined from the total p_T of tracks located within a cone $\Delta\mathcal{R} = 0.5$ around the muon. These requirements ensure that the *false* lepton is either a misidentified jet or a lepton from the semi-leptonic decay of a heavy-flavor quark. Using

n_{jet}	≥ 0	0	1	≥ 2
Channel	$\epsilon_{t\bar{t} \rightarrow ZqW\bar{b}}$ (%)			
eee	1.65 ± 0.24	$(7.65 \pm 1.45) \cdot 10^{-2}$	0.57 ± 0.09	1.00 ± 0.15
$ee\mu$	1.92 ± 0.18	$(6.77 \pm 1.05) \cdot 10^{-2}$	0.58 ± 0.06	1.17 ± 0.11
$\mu\mu e$	1.23 ± 0.13	$(3.37 \pm 0.73) \cdot 10^{-2}$	0.34 ± 0.04	0.85 ± 0.10
$\mu\mu\mu$	1.48 ± 0.19	$(3.05 \pm 0.74) \cdot 10^{-2}$	0.38 ± 0.06	1.07 ± 0.15
Channel	$\epsilon_{t\bar{t} \rightarrow ZqZ\bar{q}}$ (%)			
eee	1.22 ± 0.18	$(4.69 \pm 0.68) \cdot 10^{-2}$	0.41 ± 0.06	0.76 ± 0.11
$ee\mu$	3.75 ± 0.38	$(1.07 \pm 0.11) \cdot 10^{-1}$	1.08 ± 0.11	2.56 ± 0.25
$\mu\mu e$	1.47 ± 0.16	$(3.22 \pm 0.57) \cdot 10^{-2}$	0.38 ± 0.05	1.06 ± 0.32
$\mu\mu\mu$	2.76 ± 0.36	$(3.63 \pm 0.69) \cdot 10^{-2}$	0.63 ± 0.09	2.10 ± 0.28

Table I: Final efficiencies in % including detector and kinematic acceptance as well as detector efficiencies for each decay signature as a function of jet multiplicity n_{jet} . The efficiency, ϵ , is defined assuming fully leptonic decays of the vector bosons from top quarks, as defined as in Eq. 2. The statistical and systematic uncertainties have been added in quadrature.

a sample of data events, collected using jet triggers with no lepton requirement, we measure the ratio of misidentified leptons passing two different selection criteria, *false* lepton and signal lepton, as a function of p_T in three bins, $n_{\text{jet}} = 0, 1, \text{ and } \geq 2$, where n_{jet} is the number of jets. We then select a sample of Z boson decays with at least one additional *false* lepton candidate for each final state signature. The contribution from the V +jets background is estimated by scaling the number of events in this Z +*false* lepton sample by the corresponding p_T -dependent misidentification ratio.

Initial or final state radiation in $Z\gamma$ events can mimic the signal process if the photon either converts into an e^+e^- pair or is wrongly matched with a central track mimicking an electron and the \cancel{E}_T is mismeasured. As a result the $Z\gamma$ process is a background to the final state signatures with $W \rightarrow e\nu$ decays. To estimate the contribution from this background, we model the kinematics of these events using the $Z\gamma$ NLO MC simulation [30]. We scale this result by the rate at which a photon is misidentified as an electron. This rate is obtained using a data sample of $Z \rightarrow \mu\mu$ events containing a radiated photon, as it offers an almost background-free source of photons. The invariant mass $M(\mu\mu\gamma)$ is reconstructed and required to be consistent with the Z boson mass. The $Z \rightarrow \mu\mu$ decay is chosen to avoid any ambiguity when assigning the electromagnetic shower to the final state photon candidate. As the $Z\gamma$ NLO MC does not model recoil jets, PYTHIA MC samples are used to estimate $Z\gamma$ background jet multiplicities and \cancel{E}_T . As the PYTHIA samples do not contain events with final state radiation, we find the fraction of $Z\gamma$ events in data and PYTHIA MC that pass our \cancel{E}_T cut and take the difference as a systematic uncertainty.

After all selection criteria have been applied, we observe a total of 35 candidate events and expect $31.7 \pm 0.3(\text{stat}) \pm 3.9(\text{syst})$ background events from SM processes. The statistical uncertainty is due to MC statistics while the sources of systematic uncertainties are discussed later. Table II summarizes the number of events in each n_{jet} bin. The observed number of candidate and

background events for each topology, summing over n_{jet} , are summarized in Table III. In Tables II and III and in all the following figures, we assume a B of 5%.

To achieve better separation between signal and background, we analyze the n_{jet} and H_T distributions (defined as the scalar sum of transverse momenta of all leptons, jets, and \cancel{E}_T), and the reconstructed invariant mass for the products of the decay $t \rightarrow Zq$.

The jet multiplicity in data, SM background, and in FCNC top quark decays are shown in Fig. 2.




Figure 2: Distribution of n_{jet} for data, for simulated FCNC $t\bar{t}$ signal, and for the expected background. The $ZqZq$ signal is included in the $t\bar{t}$ FCNC contribution but is expected to be small, as can be seen from Tables II and III.

FCNC $t\bar{t}$ production leads to larger jet multiplicities and also a larger H_T . This is shown in Fig. 3.

To further increase our sensitivity we reconstruct the mass of the top quark that decays via a FCNC to a Z boson and a quark ($t \rightarrow Zq$). In events with $n_{\text{jet}} = 0$, this variable is not defined. In events with one jet, we calculate the invariant mass, $m_t^{\text{reco}} \equiv M(Z, \text{jet})$, from the 4-momenta of the jet and the identified Z boson, to reconstruct m_t . For events with two or more jets, we use the jet that gives a m_t^{reco} closest to $m_t = 172.5$ GeV.

n_{jet}	0	1	≥ 2
Background	$25.66 \pm 0.28 \pm 3.26$	$5.06 \pm 0.14 \pm 0.56$	$0.92 \pm 0.08 \pm 0.09$
$t\bar{t} \rightarrow WbZq$	0.20 ± 0.03	1.80 ± 0.27	3.87 ± 0.56
$t\bar{t} \rightarrow ZqZq$	0.002 ± 0.001	0.020 ± 0.003	0.050 ± 0.007
Observed	30	4	1

Table II: Number of observed events, expected number of $t\bar{t}$ FCNC events, and number of expected background events for each n_{jet} bin with statistical and systematic uncertainties. The MC statistical uncertainty on the $t\bar{t}$ signal is negligible, and we only present the systematic uncertainties. We assume $B = 5\%$.

Source	eee	$ee\mu$	$e\mu\mu$	$\mu\mu\mu$
WZ	$6.64 \pm 0.07 \pm 1.19$	$7.51 \pm 0.08 \pm 1.11$	$4.75 \pm 0.06 \pm 0.69$	$6.10 \pm 0.07 \pm 1.00$
ZZ	$0.33 \pm 0.03 \pm 0.06$	$1.76 \pm 0.07 \pm 0.17$	$0.46 \pm 0.04 \pm 0.07$	$1.30 \pm 0.06 \pm 0.21$
$V + \text{jets}$	$0.60 \pm 0.13 \pm 0.11$	$0.40 \pm 0.18 \pm 0.17$	$0.48 \pm 0.10 \pm 0.01$	$0.22 \pm 0.05 \pm 0.03$
$Z\gamma$	$0.18 \pm 0.05 \pm 0.08$	< 0.001	$0.66 \pm 0.07 \pm 0.38$	< 0.001
$t\bar{t} \rightarrow WbWb$	$0.04 \pm 0.01 \pm 0.01$	$0.04 \pm 0.01 \pm 0.01$	$0.05 \pm 0.01 \pm 0.01$	$0.04 \pm 0.01 \pm 0.01$
Background	$7.89 \pm 0.16 \pm 1.20$	$9.71 \pm 0.21 \pm 1.14$	$6.40 \pm 0.14 \pm 0.79$	$7.66 \pm 0.11 \pm 1.02$
$t\bar{t} \rightarrow WbZq$	1.57 ± 0.22	1.73 ± 0.17	1.17 ± 0.13	1.41 ± 0.18
$t\bar{t} \rightarrow ZqZq$	0.010 ± 0.001	0.029 ± 0.003	0.011 ± 0.001	0.022 ± 0.003
Observed	8	13	9	5

Table III: Number of observed events, expected number of $t\bar{t}$ FCNC events, and number of expected background events for each final state signature with statistical and systematic uncertainties. The MC statistical uncertainty on the $t\bar{t}$ signal is negligible, and we only present the systematic uncertainties. We assume $B = 5\%$.

The m_t^{reco} distribution is shown in Fig. 4(a). In Fig. 4(b), we present a 2-dimensional distribution of m_t^{reco} and H_T . None of the observables in Figs. 2 – 4 show evidence for the presence of FCNC in the decay of $t\bar{t}$. We therefore set 95% C.L. limits on the branching ratio $B(t \rightarrow Zq)$. The limits are derived from 10 bins of the H_T distributions for $n_{\text{jet}} = 0, 1$, and ≥ 2 . For the channels with $n_{\text{jet}} = 1$ and $n_{\text{jet}} \geq 2$, we split each H_T distribution into 4 bins in m_t^{reco} , $m_t^{\text{reco}} < 120$ GeV, $120 < m_t^{\text{reco}} < 150$ GeV, $150 < m_t^{\text{reco}} < 200$ GeV, and $m_t^{\text{reco}} > 200$ GeV.

When calculating the limit on the branching ratio we consider several sources of systematic uncertainty. The systematic uncertainties for lepton-identification efficiencies are 15% (eee), 11% ($ee\mu$), 9% ($\mu\mu e$), and 12% ($\mu\mu\mu$). The systematic uncertainty assigned to the choice of PDF is 5%. In addition, we assign 9% systematic uncertainty on $\sigma_{t\bar{t}}$ [26]. This includes the dependence on the uncertainty of m_t [27]. Furthermore, m_t is changed from 172.5 GeV to 175 GeV in $t\bar{t}$ MC samples with the difference in the result taken as a systematic uncertainty. We vary the v_{tqZ} and a_{tqZ} couplings as explained before Eq. 2, resulting in a 1% systematic uncertainty on the acceptance. Due to the uncertainty on the theoretical cross sections for WZ and ZZ production, we assign a 10% [31] systematic uncertainty to each. The major sources of systematic uncertainty on the estimated V +jets contribution arise from the \cancel{E}_T requirement and the statistics in the multijet sample used to measure the lepton-misidentification rates. These effects are estimated independently for each signature and found to be between 20% and 30%. The systematic uncertainty on the $Z\gamma$ background is estimated to be 40% and 58%

for the eee and $\mu\mu e$ channels, respectively. Uncertainties on jet energy scale, jet energy resolution, jet reconstruction and identification efficiency are estimated by varying parameters within their experimental uncertainties. For $n_{\text{jet}} = 0$ the uncertainty is found to be 1%, for $n_{\text{jet}} = 1$ it is 5%, and for $n_{\text{jet}} \geq 2$ it is 20%. The measured integrated luminosity has an uncertainty of 6.1% [32].

We use a modified frequentist approach [33] where the signal confidence level CL_s , defined as the ratio of the confidence level for the signal+background hypothesis to the background-only hypothesis ($CL_s = CL_{s+b}/CL_b$), is calculated by integrating the distributions of a test statistic over the outcomes of pseudo-experiments generated according to Poisson statistics for the signal+background and background-only hypotheses. The test statistic is calculated as a joint log-likelihood ratio (LLR) obtained by summing LLR values over the bins of the H_T distributions. Systematic uncertainties are incorporated via Gaussian smearing of Poisson probabilities for signal and backgrounds in the pseudo-experiments. All correlations between signal and backgrounds are maintained. To reduce the impact of systematic uncertainties on the sensitivity of the analysis, the individual signal and background contributions are fitted to the data, by allowing a variation of the background (or signal+background) prediction, within its systematic uncertainties [34]. The likelihood is constructed via a joint Poisson probability over the number of bins in the calculation, and is a function of scaling factors for the systematic uncertainties, which are given as Gaussian constraints associated with their priors.

We determine an observed limit of $B(t \rightarrow Zq) < 3.2\%$,






Figure 3: H_T distribution of data, FCNC $t\bar{t}$ signal, and expected background for events with (a) $n_{\text{jet}} = 0$, (b) $n_{\text{jet}} = 1$, and (c) $n_{\text{jet}} \geq 2$.

with an expected limit of $< 3.8\%$ at the 95% C.L. The limits on the branching ratio are converted to limits at the 95% C.L. on the FCNC vector, v_{tqZ} , and axial vector, a_{tqZ} , couplings as defined in Eq. 1 using the relation given in [7]. This can be done for any point in the (v_{tqZ}, a_{tqZ}) parameter space and for different quark flavors (u, c) since the differences in the helicity structure of the couplings are covered as systematic uncertainties in the limit on the branching ratio. Assuming only one non-vanishing v_{tqZ} coupling ($a_{tqZ} = 0$), we derive an observed (expected) limit of $v_{tqZ} < 0.19$ (< 0.21) for





Figure 4: (a) m_t^{reco} distribution of data, FCNC $t\bar{t}$ signal, and expected background for events with $n_{\text{jet}} \geq 1$; (b) m_t^{reco} vs. H_T distribution of data, FCNC $t\bar{t}$ signal, and background for events with $n_{\text{jet}} \geq 1$.

$m_t = 172.5$ GeV. This limit also holds assuming only one non-vanishing a_{tqZ} coupling. Figure 5 shows current limits from experiments at the LEP, HERA, and Tevatron colliders as a function of the FCNC couplings $\kappa_{tu\gamma}$ (defined in Ref. [7]) and v_{tuZ} for $m_t = 175$ GeV.

In summary, we have presented a search for top quark decays via FCNC in $t\bar{t}$ events leading to final states involving three leptons, an imbalance in transverse momentum, and jets. These final states have been explored for the first time in the context of FCNC couplings. In the absence of signal, we expect a limit of $B(t \rightarrow Zq) < 3.8\%$ and set a limit of $B(t \rightarrow Zq) < 3.2\%$ at the 95% C.L. which is currently the world's best limit. This translates into an observed limit on the FCNC coupling of $v_{tqZ} < 0.19$ for $m_t = 172.5$ GeV.

We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); FASI, Rosatom and RFBR (Russia); CNPq, FAPERJ, FAPESP and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF and KOSEF (Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); STFC and the Royal




Figure 5: Upper limits at the 95% C.L. on the anomalous $\kappa_{tu\gamma}$ and v_{tuZ} couplings assuming $m_t = 175$ GeV. Both D0 and CDF limits on v_{tqZ} are scaled to the SM cross section of $\sigma_{t\bar{t}} = 6.90$ pb [26]. Anomalous axial vector couplings and couplings of the charm quark are neglected: $a_{tuZ} = v_{tcZ} = a_{tcZ} = \kappa_{tc\gamma} = 0$. The scale parameter for the anomalous dimension-5 coupling $\kappa_{tu\gamma}$ is set to $\Lambda = m_t = 175$ GeV [16]. Any dependence of the Tevatron limits on $\kappa_{tu\gamma}$ is not displayed as the change is small and at most 6% for $\kappa_{tu\gamma} = 0.5$. The domain excluded by D0 is represented by the light (blue) shaded area. The hatched area corresponds to the additional domain excluded at HERA by the H1 experiment [16]. Also shown are upper limits obtained at LEP by the L3 experiment [15] (green dashed), at HERA by the ZEUS experiment [15] (grey dashed), and at the Tevatron by the CDF experiment [18, 19] (magenta dashed). The region above or to the right of the respective lines is excluded.

Society (United Kingdom); MSMT and GACR (Czech Republic); CRC Program and NSERC (Canada); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); and CAS and CNSF (China).

-
- [1] H. Fritzsch, Phys. Lett., B **224**, 423 (1989).
[2] H. Fritzsch, Phys. Lett. B **224**, 423 (1989).
[3] K. Nakamura *et al.* (Particle Data Group), J. Phys. G **37**, 075021 (2010).
[4] J. A. Aguilar-Saavedra, Acta Phys. Polon. B **35**, 2695 (2004).
[5] F. Larios, R. Martinez and M. A. Perez, Phys. Rev. D **72**, 057504 (2005).
[6] P. M. Ferreira, R. B. Guedes and R. Santos, Phys. Rev. D **77**, 114008 (2008).
[7] T. Han and J. L. Hewett, Phys. Rev. D **60**, 074015 (1999).
[8] T. Aaltonen *et al.* [CDF Collaboration], Phys. Rev. D **79**, 011104(R) (2009).
[9] J. T. Wei, P. Chang *et al.* [BELLE Collaboration], Phys. Rev. Lett. **103**, 171801 (2009).
[10] B. Aubert *et al.* [BaBar Collaboration], Phys. Rev. D **79**, 031102(R) (2009).
[11] S. Adler *et al.* [E949 Collaboration], Phys. Rev. D **77**, 052003 (2008).
[12] V. M. Abazov *et al.* [D0 Collaboration], Phys. Rev. Lett. **100**, 101801 (2008).
[13] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999).
[14] S. Casagande *et al.*, J. High Energy Phys. **10**, 094 (2008).
[15] P. Achard *et al.* [L3 Collaboration], Phys. Lett. B **549**, 290 (2002); J. Abdallah *et al.* [DELPHI Collaboration], Phys. Lett. B **590**, 21 (2004); A. Heister *et al.* [ALEPH Collaboration], Phys. Lett. B **543**, 173 (2002); G. Abbiendi *et al.* [OPAL Collaboration], Phys. Lett. B **521**, 181 (2001).
[16] F. D. Aaron *et al.* [H1 Collaboration], Phys. Lett. B **678**, 450 (2009).
[17] S. Chekanov *et al.* [ZEUS Collaboration], Phys. Lett. B **559**, 153 (2003).
[18] F. Abe *et al.*, Phys. Rev. Lett. **80**, 2525 (1998).
[19] T. Aaltonen *et al.* [CDF Collaboration], Phys. Rev. Lett. **101**, 192002 (2008).

- [20] V. M. Abazov *et al.* [D0 Collaboration], Phys. Lett. B **693**, 81 (2010).
- [21] V. M. Abazov *et al.* [D0 Collaboration], Phys. Lett. B **695**, 67 (2011).
- [22] V. M. Abazov *et al.* [D0 Collaboration], Nucl. Instrum. Methods Phys. Res. A **565**, 463 (2006).
- [23] G. C. Blazey *et al.*, in *Proceedings of the Workshop: "QCD and Weak Boson Physics in Run II,"* edited by U. Baur, R. K. Ellis, and D. Zeppenfeld, (Fermilab, Batavia, IL, 2000) p. 47; see Sec. 3.5 for details.
- [24] T. Sjöstrand, S. Mrenna, and P. Skands, J. High Energy Phys. **05**, 026 (2006); we used V6.419 and tune A.
- [25] M. L. Mangano *et al.*, J. High Energy Phys. **07**, 1 (2003).
- [26] S. Moch and P. Uwer, Phys. Rev. D **78**, 034003 (2008).
- [27] CDF and D0 Collaborations, arXiv:1007.3178 [hep-ex].
- [28] GEANT Detector Description and Simulation Tool, CERN Program Library Long Writeup W5013.
- [29] E. Boos *et al.* [CompHEP Collaboration], Nucl. Instrum. Meth. A **534**, 250 (2004).
- [30] U. Baur and E. Berger, Phys. Rev. D **47**, 4889 (1993).
- [31] J. M. Campbell and R. K. Ellis, Phys. Rev. D **60**, 113006 (1999).
- [32] T. Andeen *et al.*, FERMILAB-TM-2365 (2007).
- [33] T. Junk, Nucl. Instrum. Methods in Phys. Res. A **434**, 435 (1999); A. Read, in *"1st Workshop on Confidence Limits,"* CERN Report No. CERN-2000-005, 2000.
- [34] W. Fisher, FERMILAB-TM-2386-E.