QCD, Strings and AdS/CFT, an Introduction (I)

Robi Peschanski
(SPhT, Saclay)

School on QCD, Low-x Physics and Diffraction,
Copanello,Calabria,Italy, July 2007

- Strong Interactions and Strings

  QCD/Strings, A 33-years-old Puzzle

- From Dual Amplitudes to Strings

  The Veneziano Amplitude and beyond

- States and Symmetries

  String Theory: a brief introduction

- Why 26 (or 10) Dimensions?

  The Origin of Extra-Dimensions

Strong Interactions and Strings

Strong Interactions History

Strings $\not\Leftrightarrow$ QCD $\Leftrightarrow$ Strings

$\boxed{1968} \Rightarrow \boxed{1974} \Rightarrow \boxed{1998} \Rightarrow \boxed{2007} \ldots$

$\boxed{1968}$



Veneziano Amplitude

$A_R(s,q^2)$



Shapiro-Virasoro Amplitude

$A_P(s,q^2)$

The Origin of String Theory





Veneziano Amplitude

$A_R(s,q^2)$

Open String

Shapiro-Virasoro Amplitude

$A_P(s,q^2)$

Closed String

$$Open\ String: \quad \boxed{B_4 = \frac{\Gamma(-\alpha(s))\Gamma(-\alpha(t))}{\Gamma(-\alpha(s)-\alpha(t))}}$$

Regge Trajectory: $\boxed{\alpha(x) = \alpha(0) + \alpha' x}$

$$Resonances: \quad \frac{\Gamma(-\alpha(s))\Gamma(-\alpha(t))}{\Gamma(-\alpha(s)-\alpha(t))} \sim \frac{Polynomial(t\ or\ s)}{(n-\alpha(s\ or\ t))}\ when\ \alpha(s\ or\ t) \to n$$

$$ReggePoles: \quad \frac{\Gamma(-\alpha(s))\Gamma(-\alpha(t))}{\Gamma(-\alpha(s)-\alpha(t))} \sim s^{\alpha(t)}\Gamma(-\alpha(t))\ when\ s \to \infty$$

What are all these states for?

String Apparatus



World Sheet

Target Space

String States

- String Oscillator Modes

$$\boxed{[a_{n\mu}, a^{\dagger}_{m\nu}] = \eta_{\mu\nu}\delta_{nm} \ ; \ [\hat{q}_\mu, \hat{p}_\nu] = i\eta_{\mu\nu} \ ; \ \textit{D–dimensional metrics } \eta_{\mu\nu} = \left\{1, -1; -1^{*(d-2)}\right\}}$$

- String Position Operator $X^\mu(\sigma=0,\tau)$

$$\boxed{X^\mu(\sigma=0,\tau)(z) = Q^{(+)}_\mu(z) + Q^{(0)}_\mu(z) + Q^{(-)}_\mu(z) \ \ ; \ \ z = e^{i\tau}}$$

$$Q^{(+)} = i\sqrt{2\alpha'}\sum_{n=1}^{\infty}\frac{a_n}{\sqrt{n}}z^{-n} \ ; \ Q^{(-)} = -i\sqrt{2\alpha'}\sum_{n=1}^{\infty}\frac{a^{\dagger}_n}{\sqrt{n}}z^{n} \ ; \ Q^{(0)} = \hat{q} - 2i\alpha'\hat{p}\log z$$

- Relation with the Veneziano Amplitude

$$\boxed{B_4 = \int_0^1 dz \boxed{z^{-1-\alpha(s)}(1-z)^{-1-\alpha(t)}} = \frac{\Gamma(-\alpha(s))\Gamma(-\alpha(t))}{\Gamma(-\alpha(s)-\alpha(t))}}$$

$$(2\pi)^d\delta^{(d)}(\sum_{i=1}^{N}p_i)\, B_N = \int_0^1 \prod_{i=3}^{N-1} dz_i \prod_{i=2}^{N-1}\theta(z_i - z_{i+1}) \times \langle 0, p_1 | \boxed{\prod_{i=2}^{N-1} : e^{ip_i\cdot Q_i(z)} :} | 0, p_N\rangle$$

String Symmetries

- Generators of the String Conformal Symmetry

$$\boxed{[L_0, Q(z)] = -z\frac{dQ}{dz} \quad ; \quad [L_{-1}, Q(z)] = -\frac{dQ}{dz} \quad ; \quad [L_1, Q(z)] = -z^2\frac{dQ}{dz}}$$

Dilatation ; Translation ; Inversion

- Generalized Conformal Symmetry $[L_n, Q(z)] = -z^{n+1}\frac{dQ}{dz}$

$$\boxed{L_n = \sqrt{2\alpha' n}\hat{p}\cdot a_n + \sum_{m=1}^{\infty}\sqrt{m(n+m)}a_{n+m}\cdot a_m + \frac{1}{2}\sum_{m=1}^{n-1}\sqrt{m(n-m)}a_{m-n}\cdot a_m}$$

- The main property: Virasoro Algebra

$$\boxed{[L_n, L_m] = (n-m)L_{n+m} + \frac{d}{24}\, n(n^2-1)\delta_{n+m;0}}$$

- Generalized Invariance

$$[L_{\pm 1}, L_0] = \pm L_{\pm 1} \; ; \; [L_1, L_{-1}] = 2L_0$$

$$[L_n, L_m] = (n-m)L_{n+m} \; + \; \text{“}Central\ Charge\text{”}$$

WHY 26 Dimensions?

- (Covariant) Gauge Condition in QED :

$$\partial_\mu A_\mu = 0 \rightarrow q_\mu a_\mu^\dagger |0\rangle = 0 \; ; q_\mu \equiv (q, o, o, q)$$

$$\begin{array}{rcll} a_{2,3}^\dagger \; |0\rangle & = & |\phi_{2,3}\rangle & Transverse\ \gamma \quad \sum |\phi_{2,3}\rangle|^2 = 1 \\ a_0^\dagger - a_1^\dagger |0\rangle & = & |l\rangle & Longitudinal\ \gamma \quad \langle l|l\rangle = 0 \\ a_0^\dagger + a_1^\dagger |0\rangle & = & |s\rangle & Spurious\ \gamma \quad q_\mu a_\mu^\dagger \neq 0 \end{array}$$

- Covariant Gauge Conditions for Strings :

$$L_n |\phi_{string}\rangle = 0 \; for \; n > 0$$

$$\begin{array}{rcll} L_n |\phi_{string}\rangle & = & 0 & On-shell\ States \\ \langle l_{string}|l_{string}\rangle & = & 0 & Zero-Norm\ States \\ L_n |s_{string}\rangle & \neq & 0 & Spurious\ States \end{array}$$

- Decoupling of Spurious States: $|s\rangle = L_1^\dagger |\phi_1\rangle + L_2^\dagger |\phi_2\rangle$

$$\boxed{\left\{ L_2 + \frac{3}{2} L_1 L_1 \right\} |s_{string}\rangle = \sum_i |s_{string}, i\rangle + \frac{d-26}{2} \; |\phi_{string}\rangle}$$

- From: $[L_2, L_{-2}] = 4L_0 + \frac{d}{4} \Rightarrow \boxed{d=26}$
  Zero-mass states with Gauge Bosons and Graviton...

QCD/Strings: The 33-years-old Puzzle

$\boxed{1974}$

QCD Amplitudes: are they 4-D String Amplitudes?

Problems for 4-D Strings ...

- Conformal anomaly: D=26,10
- Gauge, Gravitation in the spectrum: $\alpha_{\mathcal{R}} = 1$, $\alpha_{\mathcal{P}} = 2$
- Zero mass asymptotic states
- QCD? asymptotic freedom, weak coupling,...
  ...all Standard Model physics!

...And for QCD

- QCD at Strong coupling? a String Theory?
- Strong Interactions: confinement, resonances, unitarity?
- Can we elaborate a Gauge/String Theory?

QCD, Strings and AdS/CFT, an Introduction (II)

Robi Peschanski
(SPhT, Saclay)
School on QCD, Low-x Physics and Diffraction,
Copanello,Calabria, Italy, July 2007

- Gauge/Gravity Correspondence

  Gauge Fields/String: "Duality" and "Holography"
- AdS/CFT Correspondence for S^4QCD

  Extensions beyond CFT
- A case study: AdS/CFT and QGP Hydrodynamics

  Emergence of a 5d Black Hole Geometry
- AdS/CFT and QGP Dynamics

  Quasi-Perfect Fluidity,Viscosity,Thermalization,...
- Conclusions and Prospects

  Progress and Open Problems of the QCD/String Connection

The Gauge-Gravity Correspondence

"Duality": Open String $\Leftrightarrow$ Closed String



Schomerus, 2006

$$
\begin{aligned}
Closed\ String &\Leftrightarrow 1-loop\ Open\ String \\
D-Brane\ \text{“}Universe\text{”} &\Rightarrow Open\ String\ Ending \\
Gravity &\Leftrightarrow Gauge \\
Large/Small\ Distance &\Rightarrow Gravity/Gauge\ Correspondence
\end{aligned}
$$

AdS/CFT Correspondence

J.Maldacena

1998

AdS



Gravity Source

Macroscopic

$(g_S N)^{-1} \to 0$

$AdS_5 \times S_5$ Superstring

Weak Gravity

Strong Gravity

CFT



N-Branes

Microscopic

$g_S N \to 0$

SU(N) Gauge Theory on the N-Branes

Strong Coupling

Weak Coupling

$\Longleftrightarrow$ Duality

DUALITY

- D_3-brane Solution of Supergravity: Horowitz,Strominger, 1991

$$\boxed{ds^2 = f^{-1/2}(-dt^2 + \sum_1^3 dx_i^2) + f^{1/2}(dr^2 + r^2 d\Omega_5)}$$

"Physical" Brane + Extra-Dimensions

$$f = 1 + \frac{R^4}{r^4} \; ; \; R^4 = 4\pi\alpha'^2 g_{YM}^2 N_c$$

- "Maldacena limit":

$$\frac{\alpha'(\to 0)}{r(\to 0)} \to z \; , \; R \; fixed \; \Rightarrow g_{YM}^2 N_c \to \infty$$

Strong coupling limit

$$\boxed{ds^2 = \frac{1}{R^2 z^2}(-dt^2 + \sum_{1-3} dx_i^2 + dz^2) + R^2 d\Omega_5}$$

Background Structure: $\mathrm{AdS}_5 \times S_5$ (same R^2)

HOLOGRAPHY

Exemple of $\langle Wilson\ Lines \rangle$





NON-CONFINING

CONFINING GEOMETRY

$$\boxed{\langle e^{iP\int_C \vec{A}\cdot d\vec{l}} \rangle = \int_\Sigma e^{-\frac{Area(\Sigma)}{\alpha'}} \approx e^{-\frac{Min.Area}{\alpha'}} \times Fluctuations}$$

- Conformal case : AdS_5/CFT_4

$$\langle Wilson\ Lines \rangle = e^{T*V(L)} \sim e^{T\times 1/L}$$

- Confining case : ex: Horizon at z_h (Witten 1998)

$$\langle Wilson\ Lines \rangle = e^{T*V(L)} \sim e^{T\times L(\times z_h^2)}$$

2007 A case Study: QGP formation and Relativistic Hydrodynamics



"Abstracted" from Experiments:

- Kinematic Landscape

$$\tau = \sqrt{x_0^2 - x_1^2} \; ; \; \eta = \frac{1}{2} \log \frac{x_0 + x_1}{x_0 - x_1} \; ; \; x_T = \{x_2, x_3\}$$

- QGP: (Almost) Perfect fluid behaviour (small Viscosity)
- Fast QGP Formation

Tool: Holographic Renormalization

K.Skenderis (2002)

- Using Fefferman-Graham Coordinates:

$$\boxed{ds^2 = \frac{g_{\mu\nu}(z)\ dx^\mu dx^\nu + dz^2}{z^2}}$$

- $4d \Leftrightarrow 5d$ metric:

$$\boxed{g_{\mu\nu}(z) = g^{(0)}_{\mu\nu}(= \eta_{\mu\nu}) + z^2 g^{(2)}_{\mu\nu}(= 0) + z^4 g^{(4)}_{\mu\nu}(\propto \langle T_{\mu\nu}\rangle) + \ldots}$$

$+\ldots$: derived from Einstein Eqs.

- $4d$ Constraints

$$\boxed{T^\mu{}_\mu = 0 \quad ; \quad \mathcal{D}_\nu T^{\mu\nu} = 0}$$

EMERGENCE of the 5d BLACK HOLE

Balasubramanian,de Boer,Minic; Myers; Janik,R.P.

- 4d Perfect Fluid "on the brane"

$$\boxed{\langle T_{\mu\nu}\rangle \propto g^{(4)}_{\mu\nu} = \begin{pmatrix} 3/z_0^4 = \epsilon & 0 & 0 & 0 \\ 0 & 1/z_0^4 = p_1 & 0 & 0 \\ 0 & 0 & 1/z_0^4 = p_2 & 0 \\ 0 & 0 & 0 & 1/z_0^4 = p_3 \end{pmatrix}}$$

- Holographic Renormalisation (Resummed)

$$ds^2 = -\frac{(1 - z^4/z_0^4)^2}{(1 + z^4/z_0^4)z^2}dt^2 + (1 + z^4/z_0^4)\frac{dx^2}{z^2} + \frac{dz^2}{z^2}$$

- $\Rightarrow 5d$ Black Brane with horizon at $z_0 \sim T_0^{-3}$

$$\boxed{ds^2 = -\frac{1 - \tilde{z}^4/\tilde{z}_0^4}{\tilde{z}^2}dt^2 + \frac{dx^2}{\tilde{z}^2} + \frac{1}{1 - \tilde{z}^4/\tilde{z}_0^4}\frac{d\tilde{z}^2}{\tilde{z}^2}}$$

$$z \longrightarrow \tilde{z} = z/\sqrt{1+\frac{z^4}{z_0^4}}$$

Viscosity on the light of duality

Consider a graviton that falls on this stack of N D3-branes
Will be absorbed by the D3 branes.
The process of absorption can be looked at from two different perspectives:



$=$



Absorption by D3 branes ($\sim$ viscosity) = absorption by black hole

$$\sigma_{abs}(\omega) \propto \int d^4x \, \frac{e^{i\omega t}}{\omega} \, \langle [T_{x_2x_3}(x), T_{x_2x_3}(0)] \rangle \Rightarrow \boxed{\frac{\eta}{s} \equiv \frac{\sigma_{abs}(0)/(16\pi\ G)}{A/(4\ G)} = \frac{1}{4\pi}}$$

Kovtun, Policastro, Son, Starinets (2001-...)

AdS/CFT: From Statics to Dynamics

R.Janik, RP (2005)

- Constraint Equations

$$\boxed{\begin{array}{c} \mathsf{T}^{\mu}{}_{\mu} \equiv -T_{\tau\tau} + \frac{1}{\tau^2} T_{yy} + 2T_{xx} = 0 \\ \mathsf{D}_{\nu} T^{\mu\nu} \equiv \tau \frac{d}{d\tau} T_{\tau\tau} + T_{\tau\tau} + \frac{1}{\tau^2} T_{yy} = 0 \end{array}}$$

- Boost-invariant Tensor

$$T_{\mu\nu} = \begin{pmatrix} f(\tau) & 0 & 0 & 0 \\ 0 & -\tau^3 \frac{d}{d\tau} f(\tau) - \tau^2 f(\tau) & 0 & 0 \\ 0 & 0 & f(\tau) + \frac{1}{2}\tau \frac{d}{d\tau} f(\tau) & 0 \\ 0 & 0 & 0 & \ldots \end{pmatrix}$$

- 1-parameter family of proper-time evolution

$f(\tau) \propto \tau^{-s}$: $T_{\mu\nu} t^{\mu} t^{\nu} \geq 0 \Rightarrow 0 < s < 4$

$f(\tau) \propto \tau^{-\frac{4}{3}}$: Perfect Fluid

$f(\tau) \propto \tau^{-1}$: Free streaming

$f(\tau) \propto \tau^{-0}$: "Full Anisotropy" $\epsilon = p_{\perp} = -p_L$

AdS/CFT $\Rightarrow$ Perfect Fluid at large τ

$$\mathfrak{R}^2 = R^{\mu\nu\alpha\beta} R_{\mu\nu\alpha\beta}$$



$s = \frac{4}{3} \pm .1$

A nonsingular background selects a moving Black Hole geometry corresponding to the perfect fluid at large proper-times

Dual of a Perfect fluid in S^4QCD "heavy ion experiment" : a Moving Black Hole

$$\boxed{v = \frac{z}{\tau^{1/3}}}$$

- Asymptotic metric

$$\boxed{ds^2 = \frac{1}{z^2}\left[-\frac{\left(1-\frac{e_0}{3}\,\frac{z^4}{\tau^{4/3}}\right)^2}{1+\frac{e_0}{3}\frac{z^4}{\tau^{4/3}}}d\tau^2+\left(1+\frac{e_0}{3}\frac{z^4}{\tau^{4/3}}\right)(\tau^2 dy^2+dx_\perp^2)\right]+\frac{dz^2}{z^2}}$$

- Black Hole off in the 5th dimension

$$Horizon: \;\; z_h = \left(\frac{3}{e_0}\right)^{\frac{1}{4}} \cdot \tau^{\frac{1}{3}} \; .$$

$$Temperature: \;\; T(\tau) \sim \frac{1}{z_h} \sim \tau^{-\frac{1}{3}}$$

$$Entropy: \;\; S(\tau) \sim Area \sim \tau \cdot \frac{1}{z_h^3} \sim const$$

AdS/CFT: Anisotropy at small τ

Kovchegov, Taliotis arXiv:0705.1234



Evaluation of τ_{iso}

$$Matching: \quad z_h^{late}(\tau) = \left(\frac{3}{e_0}\right)^{\frac{1}{4}} \equiv z_h^{early}(\tau) = \tau$$

$$Isotropization: \quad \tau_{iso} = \left(\frac{3N_c^2}{2\pi^2 e_0}\right)^{3/8}$$

$$Evaluation: \epsilon(\tau) = e_0\, \tau^{4/3}|_{\tau=.6} \sim 15\, GeV fermi^{-3} \Rightarrow \tau_{iso} \sim .3 fermi$$

Conclusions

In progress:

- Gauge-Gravity Correspondence
  A promising way towards QCD at strong couplings
- Results on AdS/CFT $\rightarrow$ S^4QCD Hydrodynamics
  Perfect Fluid, Viscosity, Isotropization
- Other studies
  Jet Quenching, Quark Dragging, and many others...

In outlook:

- Can we go beyond Boost Invariance?
  From Bjorken to Landau to "real" Hydrodynamics?
- Can we follow the flow from Ions to Hadrons?
  Initial and Final conditions for Hydrodynamics
- From S^4QCD to S^0QCD Hydrodynamics ?
  Can we construct the "Dual" of the QGP?

Some references...

——Strings————————-

- P. H. Frampton, *Dual Resonance Models And Superstrings, Singapore, Singapore: World Scientific* (1986);
- J. Polchinski, *String theory. Vol. 1: An introduction to the bosonic string, Phys. Lett.* **Cambridge, UK: Univ. Pr.** (1998);
  J. Polchinski, *String theory. Vol. 2: Superstring theory and beyond* **Cambridge, UK: Univ. Pr.** (1998);
- D. Lust and S. Theisen, *Lectures on string theory, Lect. Notes Phys. , Springer-Verlag* **346** (1989);

——AdS/CFT————————-

- J. Maldacena, *The Large N Limit of Superconformal Field Theories and Supergravity, Adv. Theor. Math. Phys.* **2** (1998) 231;

- S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *Gauge Theory Correlators from Non-Critical String Theory, Phys. Lett.* **B428** (1998) 105;
- E. Witten, *Anti De Sitter Space And Holography, Adv. Theor. Math. Phys.* **2** (1998) 253;
- , *Anti-de Sitter space, thermal phase transition, and confinement in gauge theories, Adv. Theor. Math. Phys.* **2** (1998) 505.
- O. Aharony, S.S. Gubser, J. Maldacena, H. Ooguri and Y. Oz, *Large N field theories, String Theory and Gravity, Phys.Rept.* **323** (2000)183.

Wilson Loops and AdS/CFT

- J. Maldacena, *Wilson loops in large N field theories, Phys. Rev. Lett.* **80** (1998) 4859;
- BlueS.-J. Rey and J. Yee, *Macroscopic strings as heavy quarks in large N gauge theory and anti-de Sitter supergravity,* hep-th/9803001;
- JJ. Sonnenschein and A. Loewy, *On the Supergravity Evaluation of Wilson Loop Correlators in Confining Theories, JHEP* **0001** (2000) 042.

- V. Schomerus, "Strings for Quantumchromodynamics," arXiv:0706.1209 [hep-ph].
- G. Policastro, D. T. Son and A. O. Starinets, "The shear viscosity of strongly coupled N = 4 supersymmetric Yang-Mills plasma," Phys. Rev. Lett. **87**, 081601 (2001);
- D. T. Son and A. O. Starinets, "Viscosity, Black Holes, and Quantum Field Theory," arXiv:0704.0240 [hep-th].;
- R.A. Janik and R. Peschanski, *P* **h** ys. Rev. D **73**, 045013 (2006) Asymptotic perfect fluid dynamics as a consequence of AdS/CFT, Phys. Rev. D73(2006) 045013
- R.A. Janik, *Viscous plasma evolution from gravity using AdS/CFT, Phys. Rev. Lett.* **98** (2007) 022302.

EXTRA SLIDES

Calculation of the Moving Duals

- Boost-Invariant 5-d F-G metric:

$$\boxed{ds^2 = \frac{-e^{a(\tau,z)}d\tau^2 + \tau^2 e^{b(\tau,z)}dy^2 + e^{c(\tau,z)}dx_\perp^2}{z^2} + \frac{dz^2}{z^2}}$$

- Scaling : $\boxed{v = \frac{z}{\tau^{s/4}}}$

$$[a(\tau,z), b(\tau,z), c(\tau,z)] = [a(v), b(v), c(v)] + \mathcal{O}\left(\frac{1}{\tau^{\#}}\right)$$

$$v(2a'(v)c'(v)+a'(v)b'(v)+2b'(v)c'(v))-6a'(v)-6b'(v)-12c'(v)+vc'(v)^2 = 0$$

$$3vc'(v)^2+vb'(v)^2+2vb''(v)+4vc''(v)-6b'(v)-12c'(v)+2vb'(v)c'(v) = 0$$

$$2vsb''(v) + 2sb'(v) + 8a'(v) - vsa'(v)b'(v) - 8b'(v) + vsb'(v)^2 +$$
$$4vsc''(v) + 4sc'(v) - 2vsa'(v)c'(v) + 2vsc'(v)^2 = 0 \ .$$

- Asymptotic Solution

$$\begin{aligned} a(v) &= A(v) - 2m(v) \\ b(v) &= A(v) + (2s-2)m(v) \\ c(v) &= A(v) + (2-s)m(v) \end{aligned}$$

Free Streaming and general cases

$$\boxed{v = \frac{z}{\tau^{1/4}}}$$

- Asymptotic metric

$$ds^2 = \frac{\left(-(1+\frac{v^4}{\sqrt{8}})^{\frac{1-2\sqrt{2}}{2}} (1-\frac{v^4}{\sqrt{8}})^{\frac{1+2\sqrt{2}}{2}} dt^2 + (1+\frac{v^4}{\sqrt{8}})^{\frac{1}{2}} (1-\frac{v^4}{\sqrt{8}})^{\frac{1}{2}} \tau^2 dy^2 + \right.}{z^2}$$

$$\frac{\left. + (1+\frac{v^4}{\sqrt{8}})^{\frac{1+\sqrt{2}}{2}} (1-\frac{v^4}{\sqrt{8}})^{\frac{1-\sqrt{2}}{2}} dx_\perp^2 \right)}{z^2} + \frac{dz^2}{z^2}$$

- True Singularities ?

  Ricci scalar:

$$R = -20 + \mathcal{O}\left(\frac{1}{\tau^{\#}}\right) \quad .$$

  Riemann tensor squared:

$$\mathfrak{R}^2 = R^{\mu\nu\alpha\beta} R_{\mu\nu\alpha\beta}$$

AdS/CFT: Selection of the Perfect Fluid

- Curvature invariant: $\boxed{\mathfrak{R}^2 = R^{\mu\nu\alpha\beta} R_{\mu\nu\alpha\beta}}$

$$\mathfrak{R}^2 = \frac{4}{\left(1-\Delta(s)^2 v^8\right)^4} \cdot \Big[10\,\Delta(s)^8 v^{32} - 88\,\Delta(s)^6 v^{24} + 42\,v^{24} s^2 \Delta(s)^4 +$$
$$+112\,v^{24}\Delta(s)^4 - 112\,v^{24}\Delta(s)^4 s + 36\,v^{20} s^3 \Delta(s)^2 - 72\,v^{20} s^2 \Delta(s)^2 +$$
$$+828\,\Delta(s)^4 v^{16} + 288\,v^{16}\Delta(s)^2 s - 288\,v^{16}\Delta(s)^2 - 108\,v^{16} s^2 \Delta(s)^2 +$$
$$-136\,v^{16} s^3 + 27\,v^{16} s^4 - 320\,v^{16} s + 160\,v^{16} + 296\,v^{16} s^2 + 36\,v^{12} s^3 +$$
$$-72\,v^{12} s^2 - 88\,\Delta(s)^2 v^8 + 42\,v^8 s^2 + 112\,v^8 - 112\,v^8 s + 10 \Big] + \mathcal{O}\left(\frac{1}{\tau^{\#}}\right)$$

- $\mathfrak{R}^2$ for $s = \frac{4}{3}$:

$$\boxed{\mathfrak{R}^2{}_{\text{perfect fluid}} = \frac{8(5w^{16} + 20w^{12} + 174w^8 + 20w^4 + 5)}{(1+w^4)^4}}$$

where $w = v/\Delta(\frac{4}{3})^{\frac{1}{4}} \equiv \sqrt[4]{3}\,v.$

AdS/CFT: Selection of the Perfect Fluid

$\mathfrak{R}^2$ for $s = \frac{4}{3} \pm .1$



Thermalization response-time of a perfect fluid

- Quasi-normal scalar modes of a static Black Hole in Fefferman-Graham coordinates

$$\boxed{\Delta\phi \equiv \frac{1}{\sqrt{-g}}\partial_i\left(\sqrt{-g}g^{ij}\partial_j\phi\right) = 0}$$

$$-\frac{1}{z^3}\frac{\left(1-\frac{z^4}{z_0^4}\right)^2}{\left(1+\frac{z^4}{z_0^4}\right)}\partial_t^2\phi(t,z)+\partial_z\left(\frac{1}{z^3}\left(1-\frac{z^8}{z_0^8}\right)\partial_z\phi(t,z)\right) = 0\,.$$

- Separation of variables $\phi(t,z) = e^{i\omega t}\phi(z)$

$$\phi''+\frac{1-\tilde{z}^2}{\tilde{z}(1-\tilde{z})(2-\tilde{z})}\,\phi'+\left(\frac{\omega}{\pi T}\right)^2\,\frac{1}{4\tilde{z}(1-\tilde{z})(2-\tilde{z})}\,\phi = 0$$

- Dominant Decay Time

$$\frac{\omega_c}{\pi T} \sim 3.1194 - 2.74667\ i$$

- A note on Viscosity *vs.* Q-n modes

Thermalization response-time of a perfect fluid

- Quasinormal scalar modes for the boost-invariant Black Hole geometry

$$\boxed{\Delta\phi \equiv \frac{1}{\sqrt{-g}}\partial_i\left(\sqrt{-g}g^{ij}\partial_j\phi\right) = 0}$$

$$\left[\partial_z \to \tau^{-\frac{1}{3}}\partial_v \ ; \ \partial_\tau \to \partial_\tau - \frac{1}{3}\tau^{-\frac{4}{3}}\partial_v\right]$$

$$-\frac{1}{v^3}\frac{(1+v^4)^2}{1-v^4}\partial_\tau^2\phi(\tau,v) + \tau^{-\frac{2}{3}}\ \partial_v\left(\frac{1}{v^3}(1-v^8)\partial_v\phi(\tau,v)\right) = 0$$

- Separation of variables $\phi(\tau,v) = f(\tau)\phi(v)$

$$\partial_\tau^2 f(\tau) = -\omega^2\tau^{-\frac{2}{3}}f(\tau) \ \Rightarrow \ f(\tau) = \sqrt{\tau}J_{\pm\frac{3}{4}}\left(\frac{3}{2}\omega\tau^{\frac{2}{3}}\right) \ \sim \ \tau^{\frac{1}{6}}\ e^{\frac{3}{2}i\omega\tau^{\frac{2}{3}}}$$

$$\boxed{\partial_v\left(\frac{1}{v^3}(1-v^8)\partial_v\phi(v)\right) + \omega^2\frac{1}{v^3}\frac{(1+v^4)^2}{1-v^4}\phi(v) = 0}$$

- Dominant Decay Proper-Time

$$\frac{\omega_c}{\pi T} \sim 3.1194 - 2.74667\ i$$

Conclusions II

- AdS/CFT and Hydrodynamics
  Construction of the "Dual" of a relativistic fluid
- $4d \to 5d$: Holographic Renormalisation
  5-d Metric from the 4-d $T_{\mu\nu}$
- 5d Physical Criterium
  Non-singular 5d Horizon $\Rightarrow$ Asymptotic
  4d perfect fluid
- Quasi-Normal mode analysis:
  The perfect fluid is thermally "very stable"

Unexpected AdS/CFT consequences for QCD at strong coupling?