



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

PHYSICS LETTERS B

Physics Letters B 622 (2005) 265–276

[www.elsevier.com/locate/physletb](http://www.elsevier.com/locate/physletb)

## Search for single top quark production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV

DØ Collaboration

V.M. Abazov<sup>al</sup>, B. Abbott<sup>bx</sup>, M. Abolins<sup>bo</sup>, B.S. Acharya<sup>af</sup>, M. Adams<sup>bb</sup>, T. Adams<sup>az</sup>,  
M. Agelou<sup>u</sup>, J.-L. Agram<sup>v</sup>, S.H. Ahn<sup>ah</sup>, M. Ahsan<sup>bi</sup>, G.D. Alexeev<sup>al</sup>, G. Alkhalaf<sup>ap</sup>,  
A. Alton<sup>bn</sup>, G. Alverson<sup>bm</sup>, G.A. Alves<sup>b</sup>, M. Anastasoie<sup>ak</sup>, T. Andeen<sup>bd</sup>,  
S. Anderson<sup>av</sup>, B. Andrieu<sup>t</sup>, Y. Arnaud<sup>q</sup>, A. Askew<sup>az</sup>, B. Åsman<sup>aq,ar</sup>,  
A.C.S. Assis Jesus<sup>c</sup>, O. Atramentov<sup>bg</sup>, C. Autermann<sup>x</sup>, C. Avila<sup>k</sup>, F. Badaud<sup>p</sup>,  
A. Baden<sup>bk</sup>, B. Baldin<sup>ba</sup>, P.W. Balm<sup>aj</sup>, S. Banerjee<sup>af</sup>, E. Barberis<sup>bm</sup>, P. Bargassa<sup>cb</sup>,  
P. Baringer<sup>bh</sup>, C. Barnes<sup>at</sup>, J. Barreto<sup>b</sup>, J.F. Bartlett<sup>ba</sup>, U. Bassler<sup>t</sup>, D. Bauer<sup>be</sup>,  
A. Bean<sup>bh</sup>, S. Beauceron<sup>t</sup>, M. Begalli<sup>c</sup>, M. Begel<sup>bt</sup>, A. Bellavance<sup>bq</sup>, S.B. Beri<sup>ad</sup>,  
G. Bernardi<sup>t</sup>, R. Bernhard<sup>ba,1</sup>, I. Bertram<sup>as</sup>, M. Besançon<sup>u</sup>, R. Beuselinck<sup>at</sup>,  
V.A. Bezzubov<sup>ao</sup>, P.C. Bhat<sup>ba</sup>, V. Bhatnagar<sup>ad</sup>, M. Binder<sup>ab</sup>, C. Biscarat<sup>as</sup>,  
K.M. Black<sup>bl</sup>, I. Blackler<sup>at</sup>, G. Blazey<sup>bc</sup>, F. Blekman<sup>at</sup>, S. Blessing<sup>az</sup>, D. Bloch<sup>v</sup>,  
U. Blumenschein<sup>z</sup>, A. Boehnlein<sup>ba</sup>, O. Boeriu<sup>bf</sup>, T.A. Bolton<sup>bi</sup>, F. Borchering<sup>ba</sup>,  
G. Borissov<sup>as</sup>, K. Bos<sup>aj</sup>, E.E. Boos<sup>an</sup>, T. Bose<sup>bs</sup>, A. Brandt<sup>bz</sup>, R. Brock<sup>bo</sup>,  
G. Brooijmans<sup>bs</sup>, A. Bross<sup>ba</sup>, N.J. Buchanan<sup>az</sup>, D. Buchholz<sup>bd</sup>, M. Buehler<sup>bb</sup>,  
V. Buescher<sup>z</sup>, V. Bunichev<sup>an</sup>, S. Burdin<sup>ba</sup>, S. Burke<sup>av</sup>, T.H. Burnett<sup>cd</sup>, E. Busato<sup>t</sup>,  
C.P. Buszello<sup>at</sup>, J.M. Butler<sup>bl</sup>, J. Cammin<sup>bt</sup>, S. Caron<sup>aj</sup>, W. Carvalho<sup>c</sup>, B.C.K. Casey<sup>by</sup>,  
N.M. Cason<sup>bf</sup>, H. Castilla-Valdez<sup>ai</sup>, S. Chakrabarti<sup>af</sup>, D. Chakraborty<sup>bc</sup>, K.M. Chan<sup>bt</sup>,  
A. Chandra<sup>af</sup>, D. Chapin<sup>by</sup>, F. Charles<sup>v</sup>, E. Cheu<sup>av</sup>, D.K. Cho<sup>bl</sup>, S. Choi<sup>ay</sup>,  
B. Choudhary<sup>ae</sup>, T. Christiansen<sup>ab</sup>, L. Christofek<sup>bh</sup>, D. Claes<sup>bq</sup>, B. Clément<sup>v</sup>,  
C. Clément<sup>aq,ar</sup>, Y. Coadou<sup>e,f,g,h</sup>, M. Cooke<sup>cb</sup>, W.E. Cooper<sup>ba</sup>, D. Coppage<sup>bh</sup>,  
M. Corcoran<sup>cb</sup>, A. Cothenet<sup>r</sup>, M.-C. Cousinou<sup>r</sup>, B. Cox<sup>au</sup>, S. Crépe-Renaudin<sup>q</sup>,  
D. Cutts<sup>by</sup>, H. da Motta<sup>b</sup>, M. Das<sup>bj</sup>, B. Davies<sup>as</sup>, G. Davies<sup>at</sup>, G.A. Davis<sup>bd</sup>, K. De<sup>bz</sup>,  
P. de Jong<sup>aj</sup>, S.J. de Jong<sup>ak</sup>, E. De La Cruz-Burelo<sup>bn</sup>, C. De Oliveira Martins<sup>c</sup>,  
S. Dean<sup>au</sup>, J.D. Degenhardt<sup>bn</sup>, F. Déliot<sup>u</sup>, M. Demarteau<sup>ba</sup>, R. Demina<sup>bt</sup>, P. Demine<sup>u</sup>,  
D. Denisov<sup>ba</sup>, S.P. Denisov<sup>ao</sup>, S. Desai<sup>bu</sup>, H.T. Diehl<sup>ba</sup>, M. Diesburg<sup>ba</sup>, M. Doidge<sup>as</sup>,  
H. Dong<sup>bu</sup>, S. Doulas<sup>bm</sup>, L.V. Dudko<sup>an</sup>, L. Duflot<sup>s</sup>, S.R. Dugad<sup>af</sup>, A. Duperrin<sup>r</sup>,

J. Dyer<sup>bo</sup>, A. Dyshkant<sup>bc</sup>, M. Eads<sup>bc</sup>, D. Edmunds<sup>bo</sup>, T. Edwards<sup>au</sup>, J. Ellison<sup>ay</sup>,  
 J. Elmsheuser<sup>ab</sup>, V.D. Elvira<sup>ba</sup>, S. Eno<sup>bk</sup>, P. Ermolov<sup>an</sup>, O.V. Eroshin<sup>ao</sup>, J. Estrada<sup>ba</sup>,  
 H. Evans<sup>bs</sup>, A. Evdokimov<sup>am</sup>, V.N. Evdokimov<sup>ao</sup>, J. Fast<sup>ba</sup>, S.N. Fatakia<sup>bl</sup>,  
 L. Feligioni<sup>bl</sup>, A.V. Ferapontov<sup>ao</sup>, T. Ferbel<sup>bt</sup>, F. Fiedler<sup>ab</sup>, F. Filthaut<sup>ak</sup>, W. Fisher<sup>br</sup>,  
 H.E. Fisk<sup>ba</sup>, I. Fleck<sup>z</sup>, M. Fortner<sup>bc</sup>, H. Fox<sup>z</sup>, S. Fu<sup>ba</sup>, S. Fuess<sup>ba</sup>, T. Gadfort<sup>cd</sup>,  
 C.F. Galea<sup>ak</sup>, E. Gallas<sup>ba</sup>, E. Galyaev<sup>bf</sup>, C. Garcia<sup>bt</sup>, A. Garcia-Bellido<sup>cd</sup>, J. Gardner<sup>bh</sup>,  
 V. Gavrilov<sup>am</sup>, A. Gay<sup>v</sup>, P. Gay<sup>p</sup>, D. Gelé<sup>v</sup>, R. Gelhaus<sup>ay</sup>, K. Genser<sup>ba</sup>, C.E. Gerber<sup>bb</sup>,  
 Y. Gershtein<sup>az</sup>, D. Gillberg<sup>e.f.g.h</sup>, G. Ginther<sup>bt</sup>, T. Golling<sup>y</sup>, N. Gollub<sup>aq.ar</sup>, B. Gómez<sup>k</sup>,  
 K. Gounder<sup>ba</sup>, A. Goussiou<sup>bf</sup>, P.D. Grannis<sup>bu</sup>, S. Greder<sup>c</sup>, H. Greenlee<sup>ba</sup>,  
 Z.D. Greenwood<sup>bj</sup>, E.M. Gregores<sup>d</sup>, Ph. Gris<sup>p</sup>, J.-F. Grivaz<sup>s</sup>, L. Groer<sup>bs</sup>,  
 S. Grünendahl<sup>ba</sup>, M.W. Grünewald<sup>ag</sup>, S.N. Gurzhiev<sup>ao</sup>, G. Gutierrez<sup>ba</sup>, P. Gutierrez<sup>bx</sup>,  
 A. Haas<sup>bs</sup>, N.J. Hadley<sup>bk</sup>, S. Hagopian<sup>az</sup>, I. Hall<sup>bx</sup>, R.E. Hall<sup>ax</sup>, C. Han<sup>bn</sup>, L. Han<sup>j</sup>,  
 K. Hanagaki<sup>ba</sup>, K. Harder<sup>bi</sup>, A. Harel<sup>ac</sup>, R. Harrington<sup>bm</sup>, J.M. Hauptman<sup>bg</sup>,  
 R. Hauser<sup>bo</sup>, J. Hays<sup>bd</sup>, T. Hebbeker<sup>x</sup>, D. Hedin<sup>bc</sup>, J.M. Heinmiller<sup>bb</sup>, A.P. Heinson<sup>ay</sup>,  
 U. Heintz<sup>bl</sup>, C. Hensel<sup>bh</sup>, G. Hesketh<sup>bm</sup>, M.D. Hildreth<sup>bf</sup>, R. Hirosky<sup>cc</sup>, J.D. Hobbs<sup>bu</sup>,  
 B. Hoeneisen<sup>o</sup>, M. Hohlfeld<sup>aa</sup>, S.J. Hong<sup>ah</sup>, R. Hooper<sup>by</sup>, P. Houben<sup>aj</sup>, Y. Hu<sup>bu</sup>,  
 J. Huang<sup>bc</sup>, V. Hynek<sup>l</sup>, I. Iashvili<sup>ay</sup>, R. Illingworth<sup>ba</sup>, A.S. Ito<sup>ba</sup>, S. Jabeen<sup>bh</sup>,  
 M. Jaffré<sup>s</sup>, S. Jain<sup>bx</sup>, V. Jain<sup>bv</sup>, K. Jakobs<sup>z</sup>, A. Jenkins<sup>at</sup>, R. Jesik<sup>at</sup>, K. Johns<sup>av</sup>,  
 M. Johnson<sup>ba</sup>, A. Jonckheere<sup>ba</sup>, P. Jonsson<sup>at</sup>, A. Juste<sup>ba</sup>, D. Käfer<sup>x</sup>, S. Kahn<sup>bv</sup>,  
 E. Kajfasz<sup>r</sup>, A.M. Kalinin<sup>al</sup>, J. Kalk<sup>bo</sup>, D. Karmanov<sup>an</sup>, J. Kasper<sup>bl</sup>, D. Kau<sup>az</sup>,  
 R. Kaur<sup>ad</sup>, R. Kehoe<sup>ca</sup>, S. Kermiche<sup>r</sup>, S. Kesisoglou<sup>by</sup>, A. Khanov<sup>bt</sup>, A. Kharchilava<sup>bf</sup>,  
 Y.M. Kharzhev<sup>al</sup>, H. Kim<sup>bz</sup>, T.J. Kim<sup>ah</sup>, B. Klima<sup>ba</sup>, J.M. Kohli<sup>ad</sup>, J.-P. Konrath<sup>z</sup>,  
 M. Kopal<sup>bx</sup>, V.M. Korablev<sup>ao</sup>, J. Kotcher<sup>bv</sup>, B. Kothari<sup>bs</sup>, A. Koubarovsky<sup>an</sup>,  
 A.V. Kozelov<sup>ao</sup>, J. Kozminski<sup>bo</sup>, A. Kryemadhi<sup>cc</sup>, S. Krzywdzinski<sup>ba</sup>, Y. Kulik<sup>ba</sup>,  
 A. Kumar<sup>ae</sup>, S. Kunori<sup>bk</sup>, A. Kupco<sup>n</sup>, T. Kurča<sup>w</sup>, J. Kvita<sup>l</sup>, S. Lager<sup>aq.ar</sup>, N. Lahrichi<sup>u</sup>,  
 G. Landsberg<sup>by</sup>, J. Lazoflores<sup>az</sup>, A.-C. Le Bihan<sup>v</sup>, P. Lebrun<sup>w</sup>, W.M. Lee<sup>az</sup>, A. Leflat<sup>an</sup>,  
 F. Lehner<sup>ba,l</sup>, C. Leonidopoulos<sup>bs</sup>, J. Leveque<sup>av</sup>, P. Lewis<sup>at</sup>, J. Li<sup>bz</sup>, Q.Z. Li<sup>ba</sup>,  
 J.G.R. Lima<sup>bc</sup>, D. Lincoln<sup>ba</sup>, S.L. Linn<sup>az</sup>, J. Linnemann<sup>bo</sup>, V.V. Lipaev<sup>ao</sup>, R. Lipton<sup>ba</sup>,  
 L. Lobo<sup>at</sup>, A. Lobodenko<sup>ap</sup>, M. Lokajicek<sup>n</sup>, A. Lounis<sup>v</sup>, P. Love<sup>as</sup>, H.J. Lubatti<sup>cd</sup>,  
 L. Lueking<sup>ba</sup>, M. Lynker<sup>bf</sup>, A.L. Lyon<sup>ba</sup>, A.K.A. Maciel<sup>bc</sup>, R.J. Madaras<sup>aw</sup>, P. Mättig<sup>ac</sup>,  
 C. Magass<sup>x</sup>, A. Magerkurth<sup>bn</sup>, A.-M. Magnan<sup>q</sup>, N. Makovec<sup>s</sup>, P.K. Mal<sup>af</sup>,  
 H.B. Malbouisson<sup>c</sup>, S. Malik<sup>bq</sup>, V.L. Malyshev<sup>al</sup>, H.S. Mao<sup>i</sup>, Y. Maravin<sup>ba</sup>,  
 M. Martens<sup>ba</sup>, S.E.K. Mattingly<sup>by</sup>, A.A. Mayorov<sup>ao</sup>, R. McCarthy<sup>bu</sup>, R. McCroskey<sup>av</sup>,  
 D. Meder<sup>aa</sup>, A. Melnitchouk<sup>bp</sup>, A. Mendes<sup>r</sup>, M. Merkin<sup>an</sup>, K.W. Merritt<sup>ba</sup>, A. Meyer<sup>x</sup>,  
 J. Meyer<sup>y</sup>, M. Michaut<sup>u</sup>, H. Miettinen<sup>cb</sup>, J. Mitrevski<sup>bs</sup>, J. Molina<sup>c</sup>, N.K. Mondal<sup>af</sup>,  
 R.W. Moore<sup>e.f.g.h</sup>, T. Moulik<sup>bh</sup>, G.S. Muanza<sup>w</sup>, M. Mulders<sup>ba</sup>, L. Mundim<sup>c</sup>,  
 Y.D. Mutaf<sup>bu</sup>, E. Nagy<sup>r</sup>, M. Narain<sup>bl</sup>, N.A. Naumann<sup>ak</sup>, H.A. Neal<sup>bn</sup>, J.P. Negret<sup>k</sup>,  
 S. Nelson<sup>az</sup>, P. Neustroev<sup>ap</sup>, C. Noeding<sup>z</sup>, A. Nomerotski<sup>ba</sup>, S.F. Novaes<sup>d</sup>,

T. Nunnemann<sup>ab</sup>, E. Nurse<sup>au</sup>, V. O’Dell<sup>ba</sup>, D.C. O’Neil<sup>e,f,g,h</sup>, V. Oguri<sup>c</sup>, N. Oliveira<sup>c</sup>,  
 N. Oshima<sup>ba</sup>, G.J. Otero y Garzón<sup>bb</sup>, P. Padley<sup>cb</sup>, N. Parashar<sup>bj</sup>, S.K. Park<sup>ah</sup>,  
 J. Parsons<sup>bs</sup>, R. Partridge<sup>by</sup>, N. Parua<sup>bu</sup>, A. Patwa<sup>bv</sup>, G. Pawloski<sup>cb</sup>, P.M. Perea<sup>ay</sup>,  
 E. Perez<sup>u</sup>, P. Pétroff<sup>s</sup>, M. Petteni<sup>at</sup>, R. Piegai<sup>a</sup>, M.-A. Pleier<sup>bt</sup>,  
 P.L.M. Podesta-Lerma<sup>ai</sup>, V.M. Podstavkov<sup>ba</sup>, Y. Pogorelov<sup>bf</sup>, M.-E. Pol<sup>b</sup>, A. Pompoš<sup>bx</sup>,  
 B.G. Pope<sup>bo</sup>, W.L. Prado da Silva<sup>c</sup>, H.B. Prosper<sup>az</sup>, S. Protopopescu<sup>bv</sup>, J. Qian<sup>bn</sup>,  
 A. Quadt<sup>y</sup>, B. Quinn<sup>bp</sup>, K.J. Rani<sup>af</sup>, K. Ranjan<sup>ae</sup>, P.A. Rapidis<sup>ba</sup>, P.N. Ratoff<sup>as</sup>,  
 S. Reucroft<sup>bm</sup>, M. Rijssenbeek<sup>bu</sup>, I. Ripp-Baudot<sup>v</sup>, F. Rizatdinova<sup>bi</sup>, S. Robinson<sup>at</sup>,  
 R.F. Rodrigues<sup>c</sup>, C. Royon<sup>u</sup>, P. Rubinov<sup>ba</sup>, R. Ruchti<sup>bf</sup>, V.I. Rud<sup>an</sup>, G. Sajot<sup>q</sup>,  
 A. Sánchez-Hernández<sup>ai</sup>, M.P. Sanders<sup>bk</sup>, A. Santoro<sup>c</sup>, G. Savage<sup>ba</sup>, L. Sawyer<sup>bj</sup>,  
 T. Scanlon<sup>at</sup>, D. Schaile<sup>ab</sup>, R.D. Schamberger<sup>bu</sup>, H. Schellman<sup>bd</sup>, P. Schieferdecker<sup>ab</sup>,  
 C. Schmitt<sup>ac</sup>, C. Schwanenberger<sup>y</sup>, A. Schwartzman<sup>br</sup>, R. Schwienhorst<sup>bo</sup>,  
 S. Sengupta<sup>az</sup>, H. Severini<sup>bx</sup>, E. Shabalina<sup>bb</sup>, M. Shamim<sup>bi</sup>, V. Shary<sup>u</sup>,  
 A.A. Shchukin<sup>ao</sup>, W.D. Shephard<sup>bf</sup>, R.K. Shivpuri<sup>ae</sup>, D. Shpakov<sup>bm</sup>, R.A. Sidwell<sup>bi</sup>,  
 V. Simak<sup>m</sup>, V. Sirotenko<sup>ba</sup>, P. Skubic<sup>bx</sup>, P. Slattery<sup>bt</sup>, R.P. Smith<sup>ba</sup>, K. Smolek<sup>m</sup>,  
 G.R. Snow<sup>bq</sup>, J. Snow<sup>bw</sup>, S. Snyder<sup>bv</sup>, S. Söldner-Rembold<sup>au</sup>, X. Song<sup>bc</sup>,  
 L. Sonnenschein<sup>t</sup>, A. Sopczak<sup>as</sup>, M. Sosebee<sup>bz</sup>, K. Soustruznik<sup>l</sup>, M. Souza<sup>b</sup>,  
 B. Spurlock<sup>bz</sup>, N.R. Stanton<sup>bi</sup>, J. Stark<sup>q</sup>, J. Steele<sup>bj</sup>, K. Stevenson<sup>be</sup>, V. Stolin<sup>am</sup>,  
 A. Stone<sup>bb</sup>, D.A. Stoyanova<sup>ao</sup>, J. Strandberg<sup>aq,ar</sup>, M.A. Strang<sup>bz</sup>, M. Strauss<sup>bx</sup>,  
 R. Ströhmer<sup>ab</sup>, D. Strom<sup>bd</sup>, M. Strovink<sup>aw</sup>, L. Stutte<sup>ba</sup>, S. Sumowidagdo<sup>az</sup>,  
 A. Sznajder<sup>c</sup>, M. Talby<sup>r</sup>, P. Tamburello<sup>av</sup>, W. Taylor<sup>e,f,g,h</sup>, P. Telford<sup>au</sup>, J. Temple<sup>av</sup>,  
 M. Tomoto<sup>ba</sup>, T. Toole<sup>bk</sup>, J. Torborg<sup>bf</sup>, S. Towers<sup>bu</sup>, T. Trefzger<sup>aa</sup>, S. Trincz-Duvoid<sup>t</sup>,  
 B. Tuchming<sup>u</sup>, C. Tully<sup>br</sup>, A.S. Turcot<sup>au</sup>, P.M. Tuts<sup>bs</sup>, L. Uvarov<sup>ap</sup>, S. Uvarov<sup>ap</sup>,  
 S. Uzunyan<sup>bc</sup>, B. Vachon<sup>e,f,g,h</sup>, P.J. van den Berg<sup>aj</sup>, R. Van Kooten<sup>be</sup>,  
 W.M. van Leeuwen<sup>aj</sup>, N. Varelas<sup>bb</sup>, E.W. Varnes<sup>av</sup>, A. Vartapetian<sup>bz</sup>, I.A. Vasilyev<sup>ao</sup>,  
 M. Vaupel<sup>ac</sup>, P. Verdier<sup>w</sup>, L.S. Vertogradov<sup>al</sup>, M. Verzocchi<sup>bk</sup>, F. Villeneuve-Segui<sup>at</sup>,  
 J.-R. Vlimant<sup>t</sup>, E. Von Toerne<sup>bi</sup>, M. Vreeswijk<sup>aj</sup>, T. Vu Anh<sup>s</sup>, H.D. Wahl<sup>az</sup>, L. Wang<sup>bk</sup>,  
 J. Warchol<sup>bf</sup>, G. Watts<sup>cd</sup>, M. Wayne<sup>bf</sup>, M. Weber<sup>ba</sup>, H. Weerts<sup>bo</sup>, N. Wermes<sup>y</sup>,  
 A. White<sup>bz</sup>, V. White<sup>ba</sup>, D. Wicke<sup>ba</sup>, D.A. Wijngaarden<sup>ak</sup>, G.W. Wilson<sup>bh</sup>,  
 S.J. Wimpenny<sup>ay</sup>, J. Wittlin<sup>bl</sup>, M. Wobisch<sup>ba</sup>, J. Womersley<sup>ba</sup>, D.R. Wood<sup>bm</sup>,  
 T.R. Wyatt<sup>au</sup>, Q. Xu<sup>bn</sup>, N. Xuan<sup>bf</sup>, S. Yacoub<sup>bd</sup>, R. Yamada<sup>ba</sup>, M. Yan<sup>bk</sup>, T. Yasuda<sup>ba</sup>,  
 Y.A. Yatsunenko<sup>al</sup>, Y. Yen<sup>ac</sup>, K. Yip<sup>bv</sup>, H.D. Yoo<sup>by</sup>, S.W. Youn<sup>bd</sup>, J. Yu<sup>bz</sup>,  
 A. Yurkewicz<sup>bu</sup>, A. Zabi<sup>s</sup>, A. Zatserklyaniy<sup>bc</sup>, M. Zdrazil<sup>bu</sup>, C. Zeitnitz<sup>aa</sup>, D. Zhang<sup>ba</sup>,  
 X. Zhang<sup>bx</sup>, T. Zhao<sup>cd</sup>, Z. Zhao<sup>bn</sup>, B. Zhou<sup>bn</sup>, J. Zhu<sup>bu</sup>, M. Zielinski<sup>bt</sup>, D. Zieminska<sup>be</sup>,  
 A. Zieminski<sup>be</sup>, R. Zitoun<sup>bu</sup>, V. Zutshi<sup>bc</sup>, E.G. Zverev<sup>an</sup>

<sup>a</sup> Universidad de Buenos Aires, Buenos Aires, Argentina<sup>b</sup> LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil<sup>c</sup> Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil<sup>d</sup> Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil

- <sup>c</sup> University of Alberta, Edmonton, AB, Canada  
<sup>f</sup> Simon Fraser University, Burnaby, BC, Canada  
<sup>g</sup> York University, Toronto, ON, Canada  
<sup>h</sup> McGill University, Montreal, Quebec, Canada  
<sup>i</sup> Institute of High Energy Physics, Beijing, People's Republic of China  
<sup>j</sup> University of Science and Technology of China, Hefei, People's Republic of China  
<sup>k</sup> Universidad de los Andes, Bogotá, Colombia  
<sup>l</sup> Center for Particle Physics, Charles University, Prague, Czech Republic  
<sup>m</sup> Czech Technical University, Prague, Czech Republic  
<sup>n</sup> Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic  
<sup>o</sup> Universidad San Francisco de Quito, Quito, Ecuador  
<sup>p</sup> Laboratoire de Physique Corpusculaire, IN2P3-CNRS, Université Blaise Pascal, Clermont-Ferrand, France  
<sup>q</sup> Laboratoire de Physique Subatomique et de Cosmologie, IN2P3-CNRS, Université de Grenoble 1, Grenoble, France  
<sup>r</sup> CPPM, IN2P3-CNRS, Université de la Méditerranée, Marseille, France  
<sup>s</sup> IN2P3-CNRS, Laboratoire de l'Accélérateur Linéaire, Orsay, France  
<sup>t</sup> LPNHE, IN2P3-CNRS, Universités Paris VI and VII, Paris, France  
<sup>u</sup> DAPNIA/Service de Physique des Particules, CEA, Saclay, France  
<sup>v</sup> IReS, IN2P3-CNRS, Université Louis Pasteur, Strasbourg, France, and Université de Haute Alsace, Mulhouse, France  
<sup>w</sup> Institut de Physique Nucléaire de Lyon, IN2P3-CNRS, Université Claude Bernard, Villeurbanne, France  
<sup>x</sup> III. Physikalisches Institut A, RWTH Aachen, Aachen, Germany  
<sup>y</sup> Physikalisches Institut, Universität Bonn, Bonn, Germany  
<sup>z</sup> Physikalisches Institut, Universität Freiburg, Freiburg, Germany  
<sup>aa</sup> Institut für Physik, Universität Mainz, Mainz, Germany  
<sup>ab</sup> Ludwig-Maximilians-Universität München, München, Germany  
<sup>ac</sup> Fachbereich Physik, University of Wuppertal, Wuppertal, Germany  
<sup>ad</sup> Panjab University, Chandigarh, India  
<sup>ae</sup> Delhi University, Delhi, India  
<sup>af</sup> Tata Institute of Fundamental Research, Mumbai, India  
<sup>ag</sup> University College Dublin, Dublin, Ireland  
<sup>ah</sup> Korea Detector Laboratory, Korea University, Seoul, South Korea  
<sup>ai</sup> CINVESTAV, Mexico City, Mexico  
<sup>aj</sup> FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands  
<sup>ak</sup> Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands  
<sup>al</sup> Joint Institute for Nuclear Research, Dubna, Russia  
<sup>am</sup> Institute for Theoretical and Experimental Physics, Moscow, Russia  
<sup>an</sup> Moscow State University, Moscow, Russia  
<sup>ao</sup> Institute for High Energy Physics, Protvino, Russia  
<sup>ap</sup> Petersburg Nuclear Physics Institute, St. Petersburg, Russia  
<sup>aq</sup> Lund University, Lund, Sweden, Royal Institute of Technology and Stockholm University, Stockholm, Sweden  
<sup>ar</sup> Uppsala University, Uppsala, Sweden  
<sup>as</sup> Lancaster University, Lancaster, United Kingdom  
<sup>at</sup> Imperial College, London, United Kingdom  
<sup>au</sup> University of Manchester, Manchester, United Kingdom  
<sup>av</sup> University of Arizona, Tucson, AZ 85721, USA  
<sup>aw</sup> Lawrence Berkeley National Laboratory and University of California, Berkeley, CA 94720, USA  
<sup>ax</sup> California State University, Fresno, CA 93740, USA  
<sup>ay</sup> University of California, Riverside, CA 92521, USA  
<sup>az</sup> Florida State University, Tallahassee, FL 32306, USA  
<sup>ba</sup> Fermi National Accelerator Laboratory, Batavia, IL 60510, USA  
<sup>bb</sup> University of Illinois at Chicago, Chicago, IL 60607, USA  
<sup>bc</sup> Northern Illinois University, DeKalb, IL 60115, USA  
<sup>bd</sup> Northwestern University, Evanston, IL 60208, USA  
<sup>be</sup> Indiana University, Bloomington, IN 47405, USA  
<sup>bf</sup> University of Notre Dame, Notre Dame, IN 46556, USA  
<sup>bg</sup> Iowa State University, Ames, IA 50011, USA  
<sup>bh</sup> University of Kansas, Lawrence, KA 66045, USA  
<sup>bi</sup> Kansas State University, Manhattan, KA 66506, USA

- <sup>bj</sup> Louisiana Tech University, Ruston, LA 71272, USA  
<sup>bk</sup> University of Maryland, College Park, MD 20742, USA  
<sup>bl</sup> Boston University, Boston, MA 02215, USA  
<sup>bm</sup> Northeastern University, Boston, MA 02115, USA  
<sup>bn</sup> University of Michigan, Ann Arbor, MI 48109, USA  
<sup>bo</sup> Michigan State University, East Lansing, MI 48824, USA  
<sup>bp</sup> University of Mississippi, University, MS 38677, USA  
<sup>bq</sup> University of Nebraska, Lincoln, NE 68588, USA  
<sup>br</sup> Princeton University, Princeton, NJ 08544, USA  
<sup>bs</sup> Columbia University, New York, NY 10027, USA  
<sup>bt</sup> University of Rochester, Rochester, NY 14627, USA  
<sup>bu</sup> State University of New York, Stony Brook, NY 11794, USA  
<sup>bv</sup> Brookhaven National Laboratory, Upton, NY 11973, USA  
<sup>bw</sup> Langston University, Langston, OK 73050, USA  
<sup>bx</sup> University of Oklahoma, Norman, OK 73019, USA  
<sup>by</sup> Brown University, Providence, RI 02912, USA  
<sup>bz</sup> University of Texas, Arlington, TX 76019, USA  
<sup>ca</sup> Southern Methodist University, Dallas, TX 75275, USA  
<sup>cb</sup> Rice University, Houston, TX 77005, USA  
<sup>cc</sup> University of Virginia, Charlottesville, VA 22901, USA  
<sup>cd</sup> University of Washington, Seattle, WA 98195, USA

Received 20 May 2005; received in revised form 1 July 2005; accepted 13 July 2005

Available online 21 July 2005

Editor: L. Rolandi

---

## Abstract

We present a search for electroweak production of single top quarks in the  $s$ -channel and  $t$ -channel using neural networks for signal–background separation. We have analyzed  $230 \text{ pb}^{-1}$  of data collected with the DØ detector at the Fermilab Tevatron Collider at a center-of-mass energy of 1.96 TeV and find no evidence for a single top quark signal. The resulting 95% confidence level upper limits on the single top quark production cross sections are 6.4 pb in the  $s$ -channel and 5.0 pb in the  $t$ -channel.

© 2005 Elsevier B.V. All rights reserved.

PACS: 14.65.Ha; 12.15.Ji; 13.85.Qk

---

## 1. Introduction

Top quark physics provides fundamental knowledge of the strong and electroweak sectors of the standard model and offers discovery potential for physics beyond the standard model. The top quark was discovered in 1995 at the Fermilab Tevatron Collider in  $t\bar{t}$  events produced through the strong interaction [1]. The standard model predicts that proton–antiproton collisions should also produce single top quarks through the electroweak interaction. Study-

ing single top quark production will provide direct measurements of the CKM matrix element  $|V_{tb}|$  and top quark polarization, and will probe possible new physics in the top quark sector [2,3].

There are two main modes of single top quark production as shown in Fig. 1: the  $s$ -channel ( $tb$ ) process  $p\bar{p} \rightarrow t\bar{b} + X$  and the  $t$ -channel ( $tqb$ ) process  $p\bar{p} \rightarrow tq\bar{b} + X$ . The production cross sections have been calculated at next-to-leading order (NLO) in the strong coupling constant [4–8], yielding  $0.88 \pm 0.14$  pb for the  $s$ -channel and  $1.98 \pm 0.30$  pb [4,5] for the  $t$ -channel, assuming a top quark mass of  $m_t = 175$  GeV. Both the DØ and CDF Collaborations have previously performed searches for single top quark production [9,10]. Recently, CDF performed a search using

---

*E-mail address:* [schwier@fnal.gov](mailto:schwier@fnal.gov) (R. Schwienhorst).

<sup>1</sup> Visitor from University of Zurich, Zurich, Switzerland.




Fig. 1. Representative Feynman diagrams for electroweak top quark production at the Tevatron Collider. This figure shows (a) the  $s$ -channel and (b) the  $t$ -channel.

160 pb<sup>-1</sup> of data and obtained upper limits of 13.6 pb ( $s$ -channel), 10.1 pb ( $t$ -channel), and 17.8 pb ( $s + t$  combined) at 95% C.L. [11].

In this Letter, we present a new search for electroweak production of single top quarks with the DØ detector. Our search focuses on the final state in which the top quark decays into a  $b$  quark and a  $W$  boson, and the  $W$  boson then decays into an electron or a muon, and a neutrino. This gives rise to an event signature with a high transverse momentum ( $p_T$ ) lepton with significant missing transverse energy ( $\cancel{E}_T$ ) from the neutrino. In addition, the  $s$ -channel has two  $b$ -quark jets, whereas the  $t$ -channel typically has one  $b$ -quark jet together with a light-quark jet. The  $\bar{b}$ -quark jet in the  $t$ -channel, as represented in Fig. 1(b), is usually emitted in the forward direction with low  $p_T$  and is often undetected [5,8]. The main backgrounds in this analysis are the  $W$  boson production in association with jets ( $W + \text{jets}$ ), top quark pair ( $t\bar{t}$ ), and multi-jet production. We use neural networks to separate the signals from the backgrounds. In the absence of any significant evidence for signal, we build a binned likelihood from the neural network outputs to set upper limits on the single top quark production cross sections.

## 2. The DØ detector

The DØ detector consists of a central tracking system, liquid-argon/uranium calorimeters, and an iron toroid muon spectrometer [12]. The central tracking system covers the detector pseudorapidity<sup>2</sup> region  $|\eta_{\text{det}}| < 2.5$ . It includes a silicon microstrip tracker

and a scintillating fiber tracker, both located within a 2 T solenoidal magnet. The calorimeters consist of a central barrel in the region  $|\eta_{\text{det}}| < 1.1$ , and two end caps extending the coverage to  $|\eta_{\text{det}}| \approx 4.2$ . The muon system outside the calorimeter consists of a layer of tracking detectors and scintillation counters before 1.8 T toroids, followed by two similar layers after the toroids. Luminosity is measured using plastic scintillator arrays located in front of the end calorimeters.

## 3. Data set and event selection

The data were recorded between August 2002 and March 2004 using a trigger that required an electromagnetic energy cluster and a jet in the calorimeter for the electron channel, or a muon and a jet for the muon channel. The integrated luminosity is 226 pb<sup>-1</sup> for the electron channel and 229 pb<sup>-1</sup> for the muon channel.

In the electron channel, we require exactly one isolated electron [13] with  $p_T > 15$  GeV and  $|\eta_{\text{det}}| < 1.1$ . In the muon channel, events are selected by requiring exactly one isolated muon [13] with  $p_T > 15$  GeV and  $|\eta_{\text{det}}| < 2.0$ . For both channels, events are also required to have  $\cancel{E}_T > 15$  GeV. Events must have from two to four jets with the leading jet  $p_T > 25$  GeV and  $|\eta_{\text{det}}| < 2.5$ , and all other jets having  $p_T > 15$  GeV and  $|\eta_{\text{det}}| < 3.4$ . Jets are defined using a cone algorithm with radius  $\mathcal{R} = 0.5$  [14]. Misreconstructed events are rejected by requiring that the direction of  $\cancel{E}_T$  is not aligned or anti-aligned in azimuth with the lepton or the jets.

The fraction of signal-like events is further enhanced through the selection of  $b$ -quark jets that are identified by reconstructing displaced vertices from long-lived particles. A displaced vertex is selected by requiring the transverse decay-length significance,  $L_{xy}/\sigma_{L_{xy}}$ , to be greater than seven, where  $L_{xy}$  is the decay length and  $\sigma_{L_{xy}}$  is the uncertainty on  $L_{xy}$ , calculated from the error matrices of the tracks and the primary vertex. A jet is considered  $b$ -tagged by this algorithm if a displaced vertex lies within a cone of radius  $\mathcal{R} = 0.5$  around the jet axis [15].

<sup>2</sup> Pseudorapidity is defined as  $\eta = -\ln(\tan \frac{\theta}{2})$ , where  $\theta$  is the polar angle with the origin at the primary vertex. Detector fiducial

regions are defined by detector pseudorapidity  $\eta_{\text{det}}$  which is calculated with the origin at the nominal center of the detector ( $z = 0$ ).



For both  $s$ -channel and  $t$ -channel searches, we separate the data into independent analysis sets based on final-state lepton flavor (electron or muon) and  $b$ -tag multiplicity. To take advantage of the different final state topologies, we separate single-tagged ( $= 1$  tag) events from double-tagged ( $\geq 2$  tags) events. In the  $t$ -channel search, we additionally require that one of the jets is not  $b$  tagged.

#### 4. Acceptances and yields

We estimate the kinematic and geometric acceptances for  $s$ -channel and  $t$ -channel single top quark production using the COMPHEP matrix element event generator [16] with  $m_t = 175$  GeV. The factorization scales are  $m_t^2$  for the  $s$ -channel samples and  $(m_t/2)^2$  for the  $t$ -channel samples. For the  $s$ -channel ( $t$ -channel) search, the  $t$ -channel ( $s$ -channel) is considered as background.

We use both Monte Carlo events and data to estimate the background yields. The  $W +$  jets and diboson ( $WW$  and  $WZ$ ) backgrounds are estimated using events generated with ALPGEN [17]. The diboson background yields are normalized to NLO cross sections computed with MCFM [18]. The total  $W +$  jets yield is normalized to the yield in data corrected for the presence of multijet,  $t\bar{t}$  and dibosons before requiring a  $b$ -tagged jet. The fraction of heavy-flavor ( $Wb\bar{b}$ ) events is obtained using the ratio of the NLO cross sections for  $W +$  jets and  $Wb\bar{b}$ , as described in Ref. [19]. This normalization to data also accounts for smaller contributions such as  $Z +$  jets events in which one of the leptons from the  $Z$  boson decay is not reconstructed.

The  $t\bar{t}$  background, consisting of the leptonic decay modes of the  $W$  boson from the top quark decay ( $\ell +$  jets and dilepton), is estimated using samples generated with ALPGEN, normalized to the cross section:  $\sigma(t\bar{t}) = 6.7 \pm 1.2$  pb [20], where the uncertainty on the top quark mass is incorporated into the cross section uncertainty.

The parton-level samples are then processed with PYTHIA [21] for hadronization, particle decays, and modeling of the underlying event. The generated events are processed through a GEANT-based [22] simulation of the DØ detector. The resulting lepton and jet

Table 1

Estimates for signal and background yields and the numbers of observed events in data after event selection for the electron and muon, single-tagged and double-tagged analysis sets combined. The  $W +$  jets yields include the diboson backgrounds. The total background for the  $s$ -channel ( $t$ -channel) search includes the  $tqb$  ( $tb$ ) yield. The quoted yield uncertainties include systematic uncertainties taking into account correlations between the different analysis channels and samples

Source	$s$ -channel search	$t$ -channel search
$tb$	$5.5 \pm 1.2$	$4.7 \pm 1.0$
$tqb$	$8.6 \pm 1.9$	$8.5 \pm 1.9$
$W +$ jets	$169.1 \pm 19.2$	$163.9 \pm 17.8$
$t\bar{t}$	$78.3 \pm 17.6$	$75.9 \pm 17.0$
Multijet	$31.4 \pm 3.3$	$31.3 \pm 3.2$
Total background	$287.4 \pm 31.4$	$275.8 \pm 31.5$
Observed events	283	271

energies are smeared to reproduce the resolutions observed in data.

The background from jets misidentified as electrons or jets resulting in isolated muons is estimated using multijet data samples that pass all event selection cuts, but fail the requirement on muon isolation or electron quality [13]. This background is normalized using a data sample dominated by multijet events, selected by requiring  $\cancel{E}_T < 15$  GeV.

The overall acceptances, including trigger and selection efficiencies, for signal events with at least one  $b$ -tagged jet are  $(2.7 \pm 0.2)\%$  in the  $s$ -channel and  $(1.9 \pm 0.2)\%$  in the  $t$ -channel. The acceptance is calculated as the fraction of events that pass the selection over all possible single top quark decays, including all leptonic and hadronic decays of the  $W$  boson. Estimates for signal and background yields and the observed numbers of events after selection are shown in Table 1.

#### 5. Neural networks analysis

After event selection, several variables are combined in neural networks to discriminate the single top quark signals from the backgrounds. The networks are composed of three layers of nodes: input, hidden, and output. For training and testing, we use the MLPFIT [23] package. Testing and training event sets are created from simulated signal and background samples. We use a technique called early stopping

Table 2

Input variables for each neural network signal–background pair. Variable descriptions can be found in the text

Variable	Description	Signal–background pairs			
		$t\bar{b}$		$tq\bar{b}$	
		$Wb\bar{b}$	$t\bar{t}$	$Wb\bar{b}$	$t\bar{t}$
<i>Individual object kinematics</i>					
$p_T(\text{jet1}_{\text{tagged}})$	Transverse momentum of the leading tagged jet	✓	✓	✓	–
$p_T(\text{jet1}_{\text{untagged}})$	Transverse momentum of the leading untagged jet	–	–	✓	✓
$p_T(\text{jet2}_{\text{untagged}})$	Transverse momentum of the second untagged jet	–	–	–	✓
$p_T(\text{jet1}_{\text{nonbest}})$	Transverse momentum of the leading nonbest jet	✓	✓	–	–
$p_T(\text{jet2}_{\text{nonbest}})$	Transverse momentum of the second nonbest jet	✓	✓	–	–
<i>Global event kinematics</i>					
$M_T(\text{jet1}, \text{jet2})$	Transverse mass of the two leading jets	✓	–	–	–
$p_T(\text{jet1}, \text{jet2})$	Transverse momentum of the two leading jets	✓	–	✓	–
$M(\text{alljets})$	Invariant mass of all jets	✓	✓	✓	✓
$H_T(\text{alljets})$	Sum of the transverse energies of all jets	–	–	✓	–
$M(\text{alljets} - \text{jet1}_{\text{tagged}})$	Invariant mass of all jets excluding the leading tagged jet	–	–	–	✓
$H(\text{alljets} - \text{jet1}_{\text{tagged}})$	Sum of the energies of all jets excluding the leading tagged jet	–	✓	–	✓
$H_T(\text{alljets} - \text{jet1}_{\text{tagged}})$	Sum of the transverse energies of all jets excluding the leading tagged jet	–	–	–	✓
$p_T(\text{alljets} - \text{jet1}_{\text{tagged}})$	Transverse momentum of all jets excluding the leading tagged jet	–	✓	–	✓
$M(\text{alljets} - \text{jet}_{\text{best}})$	Invariant mass of all jets excluding the best jet	–	✓	–	–
$H(\text{alljets} - \text{jet}_{\text{best}})$	Sum of the energies of all jets excluding the best jet	–	✓	–	–
$H_T(\text{alljets} - \text{jet}_{\text{best}})$	Sum of the transverse energies of all jets excluding the best jet	–	✓	–	–
$M(W, \text{jet1}_{\text{tagged}})$	Invariant mass of the reconstructed top quark using the leading tagged jet	✓	✓	✓	✓
$M(W, \text{jet}_{\text{best}})$	Invariant mass of the reconstructed top quark using the best jet	✓	–	–	–
$\sqrt{s}$	Invariant mass of all final state objects	✓	–	✓	✓
<i>Angular variables</i>					
$\Delta\mathcal{R}(\text{jet1}, \text{jet2})$	Angular separation between the leading two jets	✓	–	✓	–
$\eta(\text{jet1}_{\text{untagged}}) \times Q_\ell$	Pseudorapidity of the leading untagged jet $\times$ lepton charge	–	–	✓	✓
$\cos(\ell, Q_\ell \times z)_{\text{topbest}}$	Top quark spin correlation in the optimal basis for the $s$ -channel [25], reconstructing the top quark with the best jet	✓	–	–	–
$\cos(\ell, \text{jet1}_{\text{untagged}})_{\text{toptagged}}$	Top quark spin correlation in the optimal basis for the $t$ -channel [25], reconstructing the top quark with the leading tagged jet	–	–	✓	–
$\cos(\text{alljets}, \text{jet1}_{\text{tagged}})_{\text{alljets}}$	Cosine of the angle between the leading tagged jet and the alljets system in the alljets rest frame	–	–	✓	✓
$\cos(\text{alljets}, \text{jet}_{\text{nonbest}})_{\text{alljets}}$	Cosine of the angle between the leading non-best jet and the alljets system in the alljets rest frame	–	✓	–	–

[24] to determine the maximum number of epochs for training which prevents over-training. Each network is then tuned by choosing the optimal number of hidden nodes. From studies based on optimizing the expected upper limits on the single top quark production cross sections, we find that the  $s$ -channel and  $t$ -channel searches each require only two networks, corresponding to the dominant backgrounds:  $Wb\bar{b}$  and  $t\bar{t} \rightarrow \ell + \text{jets}$ .

The list of discriminating variables has been chosen based on an analysis of Feynman diagrams of signals and backgrounds [26] and on a study of single top quark production at NLO [7,8]. The input

variables to each network are selected from this list by training with different combinations of variables and choosing the combination that produces the minimum testing error and largest signal–background separation. Table 2 shows the variables used for each signal–background pair. These variables fall into three categories: individual-object kinematics, global-event kinematics, and angular correlations.

Since the input variables do not depend on the lepton flavor, the electron and muon analyses utilize the same variables. However, owing to different lepton resolutions and pseudorapidity ranges, we construct separate networks for them. Therefore, four neural






Fig. 2. Comparison of signal, background, and data for the electron and muon channels combined, requiring at least one tag, for four representative neural network input variables. Shown are (a) the transverse momentum of the leading untagged jet, (b) the invariant mass of the reconstructed top quark using the leading tagged jet, (c) the invariant mass of the final state system, and (d) the pseudorapidity of the leading untagged jet multiplied by the charge of the lepton. Signals are multiplied by ten.

networks are used for the signal–background pairs ( $tb-Wb\bar{b}$ ,  $tb-t\bar{t}$ ,  $tqb-Wb\bar{b}$ ,  $tqb-t\bar{t}$ ) for each of the electron and muon channels.

Fig. 2 shows distributions of four representative variables. We reconstruct the final state top quark from the reconstructed  $W$  boson and a jet as follows. The  $W$  boson is reconstructed from the isolated lepton and the missing transverse energy. The  $z$ -component of the neutrino momentum ( $p_z^{\nu}$ ) is calculated using a  $W$  boson mass constraint, choosing the solution with smaller  $|p_z^{\nu}|$  from the two possible solutions [27]. In the  $s$ -channel analysis, the top quark is reconstructed from the  $W$  boson and the “best” jet [9]. The best jet is defined as the jet in each event for which the invariant mass of the reconstructed  $W$  boson and the jet system is closest to  $m_t = 175$  GeV. In the  $t$ -channel analysis, the top quark is reconstructed from the  $W$  boson and the leading  $b$ -tagged jet. Using these two methods we are able to correctly identify the  $b$ -quark jet from the top quark decay in about 90% of the signal events.

Fig. 3 shows the outputs of the neural networks for the data and the expected backgrounds, as well as signals for the electron and muon channels combined. The neural network output in MLPFIT is around one for signal events and around zero for background events, but it is not constrained to the interval  $[0, 1]$ . The  $t\bar{t}$  networks separate signal and  $t\bar{t}$  backgrounds efficiently. The  $Wb\bar{b}$  networks are less efficient for the  $W + \text{jets}$  backgrounds because the event kinematics are similar between signal and background.

## 6. Systematic uncertainties

Systematic uncertainties are evaluated for the Monte Carlo signal and background samples, separately for the electron and muon channels and for each  $b$ -tag multiplicity. The most important sources of systematic uncertainty are listed in Table 3. The systematic uncertainty on the shapes of the distributions




Fig. 3. Comparison of signal, background, and data for the neural network outputs, for the electron and muon channels combined, requiring at least one tag. This figure shows (a) the  $tb-t\bar{t}$  filter, (b) the  $tqb-t\bar{t}$  filter, (c) the  $tb-Wb\bar{b}$  filter, and (d) the  $tqb-Wb\bar{b}$  filter. Signals are multiplied by ten.

Table 3

Range of systematic uncertainty values for the various Monte Carlo signal and background samples in the different analysis channels

Source of systematic uncertainty	Uncertainty range (%)
Signal and background acceptance	
$b$ -tag modeling	5–20
jet energy calibration	1–15
trigger modeling	2–7
jet fragmentation	5–7
jet identification	1–13
lepton identification	4
Background normalization	
theory cross sections	2–18
$W$ + jets flavor composition	5–16
Luminosity	6.5

is also taken into account for the contributions from  $b$ -tag modeling, jet energy calibration, jet identification, and trigger modeling. In order to evaluate the total uncertainty, we consider all sources of systematic uncertainties for all samples and their correlations. The total uncertainty on the signal acceptance for single-

tagged events is 13% for the  $s$ -channel and 15% for the  $t$ -channel, and for double-tagged events it is 24% for the  $s$ -channel and 28% for the  $t$ -channel. The total uncertainty on the background yield is 10% for the single-tagged samples and 26% for the double-tagged samples.

## 7. Cross section limits

The observed data are consistent with the background predictions for all eight analysis channels. We therefore set upper limits on the single top quark production cross section separately in the  $s$ -channel and  $t$ -channel searches using a Bayesian approach [28]. In each search, two-dimensional histograms are constructed from the  $Wb\bar{b}$  vs.  $t\bar{t}$  neural network outputs. A likelihood is built from these histograms for signal, background, and data, as a product over all channels (electron and muon, single and double tags) and bins. We assume a Poisson distribution for the observed




Fig. 4. The Bayesian posterior probability density as a function of the single top quark cross section for the  $s$ -channel and  $t$ -channel searches.

number of events in each bin and a flat prior probability for the signal cross section. The prior for the combined signal acceptance and background yields is a multivariate Gaussian with uncertainties and correlations described by a covariance matrix. Finally, we compute the posterior probability density as a function of the production cross section.

The Bayesian posterior probability densities are shown in Fig. 4 for both the  $s$ -channel and  $t$ -channel searches. The corresponding upper limits at the 95% C.L. are 6.4 pb in the  $s$ -channel and 5.0 pb in the  $t$ -channel. The sensitivity of these measurements is given by the expected upper limits obtained by setting the observed number of events to the background prediction in each bin. The expected upper limits are 4.5 pb in the  $s$ -channel search and 5.8 pb in the  $t$ -channel search.

## 8. Conclusions

No evidence is found for electroweak production of single top quarks in  $230 \text{ pb}^{-1}$  of data collected with the DØ detector at  $\sqrt{s} = 1.96 \text{ TeV}$ . The data consist of events in the electron and muon final states with at least one  $b$ -tagged jet. We build binned likelihoods from the output of neural networks to set upper limits at the 95% C.L. The measured  $s$ -channel limit is 6.4 pb and the measured  $t$ -channel limit is 5.0 pb. These upper limits represent significant improvements over pre-

viously published results [9–11] due to the larger data set as well as the use of a multivariate analysis technique together with shape information from the resulting output distributions. They approach the region of sensitivity for models of physics beyond the standard model, such as a fourth quark-generation scenario or flavor-changing neutral-currents [3].

## Acknowledgement

We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); FASI, Rosatom and RFBR (Russia); CAPES, CNPq, FAPERJ, FAPESP and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF (Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); PPARC (United Kingdom); MSMT (Czech Republic); CRC Program, CFI, NSERC and WestGrid Project (Canada); BMBF and DFG (Germany); SFI (Ireland); Research Corporation, Alexander von Humboldt Foundation, and the Marie Curie Program.

## References

- [1] CDF Collaboration, F. Abe, et al., Phys. Rev. Lett. 74 (1995) 2626; DØ Collaboration, S. Abachi, et al., Phys. Rev. Lett. 74 (1995) 2632.
- [2] D. Chakraborty, J. Konigsberg, D.L. Rainwater, Annu. Rev. Nucl. Part. Sci. 53 (2003) 301.
- [3] T. Tait, C.-P. Yuan, Phys. Rev. D 63 (2001) 014018.
- [4] B.W. Harris, et al., Phys. Rev. D 66 (2002) 054024.
- [5] Z. Sullivan, Phys. Rev. D 70 (2004) 114012.
- [6] J. Campbell, R.K. Ellis, F. Tramontano, Phys. Rev. D 70 (2004) 094012.
- [7] Q.-H. Cao, R. Schwienhorst, C.-P. Yuan, Phys. Rev. D 71 (2005) 054023.
- [8] Q.-H. Cao, R. Schwienhorst, J. Benitez, R. Brock, C.-P. Yuan, hep-ph/0504230, Phys. Rev. D, submitted for publication.
- [9] DØ Collaboration, B. Abbott, et al., Phys. Rev. D 63 (2001) 031101; DØ Collaboration, V.M. Abazov, et al., Phys. Lett. B 517 (2001) 282.
- [10] CDF Collaboration, D. Acosta, et al., Phys. Rev. D 65 (2002) 091102.
- [11] CDF Collaboration, D. Acosta, et al., Phys. Rev. D 71 (2005) 012005.

- [12] DØ Collaboration, V.M. Abazov, et al., The upgraded DØ detector, in preparation for submission to Nucl. Instrum. Methods in Phys. Res. A.
- [13] DØ Collaboration, V.M. Abazov, et al., hep-ex/0504043.
- [14] Jets are defined using the iterative, seed-based cone algorithm with radius  $\mathcal{R} = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2} = 0.5$ , including midpoints as described in G.C. Blazey et al., in: U. Baur, R.K. Ellis, D. Zeppenfeld (Eds.), Proceedings of the Workshop on QCD and Weak Boson Physics in Run II, FERMILAB-PUB-00-297, 2000, pp. 47–77.
- [15] DØ Collaboration, V.M. Abazov, et al., hep-ex/0504058.
- [16] CompHEP Collaboration, E. Boos, et al., Nucl. Instrum. Methods A 534 (2004) 250.
- [17] M.L. Mangano, et al., JHEP 0307 (2003) 001.
- [18] J.M. Campbell, R.K. Ellis, Phys. Rev. D 60 (1999) 113006.
- [19] DØ Collaboration, V.M. Abazov, et al., Phys. Rev. Lett. 93 (2004) 141801.
- [20] R. Bonciani, et al., Nucl. Phys. B 529 (1998) 424; M. Cacciari, et al., JHEP 0404 (2004) 068; N. Kidonakis, R. Vogt, Phys. Rev. D 68 (2003) 114014.
- [21] T. Sjöstrand, et al., Comput. Phys. Commun. 135 (2001) 238.
- [22] R. Brun, et al., CERN Program Library Long Writeup W 5013 (1994).
- [23] J. Schwindling, <http://schwind.home.cern.ch/schwind/MLPfit.html>.
- [24] G. Orr, K. Müller, Neural Networks: Tricks of the Trade, Springer-Verlag, Berlin, 1998, p. 55.
- [25] G. Mahlon, S. Parke, Phys. Rev. D 53 (1996) 4886; S. Parke, Y. Shadmi, Phys. Lett. B 387 (1996) 199.
- [26] E. Boos, L. Dudko, Nucl. Instrum. Methods A 502 (2003) 486.
- [27] G.L. Kane, C.P. Yuan, Phys. Rev. D 40 (1989) 2231.
- [28] I. Bertram, et al., FERMILAB-TM-2104 (2000).