

# Supersymmetry Searches at Tevatron

## Run I Results and Run II Prospects

**Jianming Qian**  
*The University of Michigan*


- **Introduction**
- **Run I searches**
- **Run II prospects**
- **Summary**




**SUSY2K, CERN, June 27, 2000**

# Tevatron Collider (Run I)

$p\bar{p}$  collider at  $\sqrt{s} = 1.8$  TeV




DØ Run I Integrated Luminosity



Both CDF and DØ recorded over  $100 \text{ pb}^{-1}$  data




# CDF and DØ Detectors (Run I)




**Tracking ( $l, b, \dots$ )**

**DØ**



**Calorimeter ( $e, \gamma, E_T, \dots$ )**

# Supersymmetry Memu



**Nature hasn't been kind to us**

**Theorists speculate, we analyze data,  
but at the end, nothing found**

## R-parity conservation

- **Jets**
- **Dilepton**
- **Trilepton**
- **Photonic signatures**

## R-parity violation

- **Dilepton**
- **Trileptons**
- **Four-lepton**



## Analysis categories

- **Thesis driven**
- **Fashion driven**  
gauge-mediation,  
large extra dimension  
.....
- **Gene searches for physics  
beyond the standard model**

## **Results are often interpreted in models**

- **minimal supersymmetric extension of SM (MSSM)**
- **minimal super-gravity models (mSUGRA)**
- **models with gauge mediated supersymmetry breaking**


# Signal Cross Sections



**Production has less dependence on susy parameters than decays**


**Squarks and gluinos dominant the production at Tevatron if kinematically accessible**

**Cross sections for scalar leptons are small**



# Background Cross Sections

The cross section for new physics is small compared with dominant Standard Model processes




**Leptons ( $e, \mu$ ) and missing  $E_T$  are the keys**

- missing  $E_T$  resolution
- lepton identification efficiency
- lepton fake rate

# Supergravity Signatures

For most of the parameter space, the lightest neutralino is the LSP and is so assumed for most of the analyses



$$\begin{aligned} \tilde{\chi}_i^0 &\rightarrow Z \tilde{\chi}_1^0 \\ \tilde{\chi}_j^\pm &\rightarrow W \tilde{\chi}_1^0 \\ \tilde{q} &\rightarrow q \tilde{\chi}_1^0 \\ \tilde{g} &\rightarrow q \tilde{q} \end{aligned}$$

$$\begin{aligned} p\bar{p} &\rightarrow \tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow WZ + E_T \Rightarrow \ell^{1,2,3} + E_T + X \\ &\tilde{\chi}_1^\pm \tilde{\chi}_1^\pm \rightarrow WW + E_T \Rightarrow \ell^{1,2} + E_T + X \end{aligned}$$


$$\begin{aligned} p\bar{p} &\rightarrow \tilde{q}\tilde{q} \rightarrow X + \tilde{\chi}_1^0 \tilde{\chi}_1^0 \Rightarrow jets + E_T \\ &\quad + \tilde{\chi}_1^\pm \tilde{\chi}_1^0 \Rightarrow jets + E_T + \ell \\ &\quad + \tilde{\chi}_1^\pm \tilde{\chi}_1^\mp \Rightarrow jets + E_T + \ell^\pm \ell^\mp \\ &\quad + \tilde{\chi}_2^0 \tilde{\chi}_1^0 \Rightarrow jets + E_T + \ell^\pm \ell^\mp \\ &\quad + \tilde{\chi}_1^\pm \tilde{\chi}_2^0 \Rightarrow jets + E_T + lll \\ &\quad + \dots \\ p\bar{p} &\rightarrow \tilde{g}\tilde{g} \rightarrow X + \tilde{\chi}_1^\pm \tilde{\chi}_1^\pm \Rightarrow jets + E_T + \ell^\pm \ell^\pm \end{aligned}$$

## Signatures:

$$p\bar{p} \rightarrow SUSY \Rightarrow E_T + \ell^n + j^m$$

# Stop and Sbottom Signatures


In many supersymmetry models, stop (and sbottom) can be significantly lighter than other squarks




$$\begin{aligned} \tilde{t}_1 &\rightarrow b + \tilde{\chi}_1^+ \rightarrow Wb + \tilde{\chi}_1^0 \\ \tilde{t}_1 &\rightarrow b + \tilde{\chi}_1^+ \rightarrow bl + \tilde{\nu} \\ \tilde{t}_1 &\rightarrow \tilde{\chi}_1^0 + c \\ \tilde{t}_1 &\rightarrow t + \tilde{\chi}_1^0 \end{aligned}$$

**Signatures:**

- 1) excess of SM top events
- 2) two acoplanar c-jets with  $mE_T$
- 3) two b-jets, two leptons and  $mE_T$



Assuming  $Br(\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0) = 100\%$   
pair production of  $\tilde{b}_1$  will yield  
two acoplanar b-jets




**Signatures:**

**two acoplanar b-jets + missing  $E_T$**

(Also expected from WH, ZH)

# Squarks and Gluinos

Squarks and gluinos can be copiously produced at Tevatron if they are light





The dominant signature for  $p\bar{p} \rightarrow \tilde{q}\tilde{q}, \tilde{q}\tilde{g}, \tilde{g}\tilde{g} + X$  production is therefore *jets* +  $\cancel{E}_T$

**DØ: Phys. Rev. Lett. 82, 29 (1999)**

- 1)  $\geq 3$  jets with  $E_T > 25$  GeV  
 $|\eta| < 1.1$  or  $1.4 < |\eta| < 3.5$
- 2)  $E_T^{j1} > 115$  GeV
- 3)  $\cancel{E}_T > 75$  GeV
- 4) topological cuts to reduce instrumental backgrounds

Final selection criteria for each  $(m_0, m_{1/2})$  were determined by choosing  $H_T$  and  $\cancel{E}_T$  thresholds that maximized the  $S/\delta B$  ratio

The background is dominated by QCD multijet events  
No apparent excess is found beyond expectation




# Stop and Sbottom from Jets+mEt

**CDF searched for events with heavy-quark jets and large mEt.**



- 1) 2 or 3 jets with  $E_T > 15$  GeV  
no jet with  $7 < E_T < 15$  GeV
- 2)  $E_T > 40$  GeV
- 3) no identified leptons with  $E_T > 10$  GeV
- 4) topological cuts
- 5) jet probability  $< 0.05$  for  $\tilde{t}_1$  searches  
 $< 0.01$  for  $\tilde{b}$  searches

**Backgrounds are dominated by  $W(\rightarrow\tau\nu)$ +jets and mismeasurements**



**Stop searches:  
11 events observed  
~15 expected**

**Sbottom searches:  
5 events observed  
~6 expected**



# Stop from b-jets, lepton and mEt



**CDF searched for  $\tilde{t}_1\tilde{t}_1$  production with  $\tilde{t}_1 \rightarrow b + \tilde{\chi}_1^+$  and  $\tilde{\chi}_1^+ \rightarrow W\tilde{\chi}_1^0$  or  $\ell\tilde{\nu}$**

**CDF: Phys. Rev. Lett. 84, 5273 (2000)**


- 1)  $\geq 2\ell$  with  $E_T > 10$  GeV, but not  $e^+e^-$ ,  $\mu^+\mu^-$
- 2)  $\geq 2$  jets with  $E_T > 12,8$  GeV,  $\geq 1$  vertex tagged
- 3)  $E_T > 25$  GeV
- 4) cleanup requirements to reduce mismeasurement backgrounds

**81 events were observed compared with  $87.3 \pm 8.8$  expected from background processes dominated by W+jets, tt, bb and fakes**

**Unbinned likelihood fit to kinematic distributions to extract potential stop signal. Results are consistent with no stop production**



# Dilepton ( $\pm\pm$ ) from Squarks and Gluinos



Like-sign dileptons are expected from

$$p\bar{p} \rightarrow \tilde{q}\tilde{q}^*, \tilde{q}\tilde{g}, \tilde{g}\tilde{g} \Rightarrow l^\pm l^\pm jj E_T + X$$

- 1)  $2l^\pm l^\pm$  with  $E_T > 11,5$  GeV
- 2)  $\geq 2$  jets with  $E_T > 15$  GeV
- 3) dilepton mass cut
- 4)  $E_T > 25$  GeV


- Backgrounds:**
- Drell-Yan
  - heavy quarks
  - dibosons

No like-sign events were observed while  $0.6 \pm 0.3$  events were expected from known processes.

19 opposite-sign events passed the similar selection.

The null results is used to constrain squark/gluino production within the SUGRA-inspired MSSM and assuming 5 degenerate squarks.

The efficiency for the signal is about 1%.



# Dilepton from mSUGRA

Squarks and gluinos can also result in dilepton final states

$$p\bar{p} \rightarrow SUSY \Rightarrow \ell\ell jj + \cancel{E}_T$$

- 1) two leptons (e,  $\mu$ ) with varying  $E_T$  cut
- 2) two or three jets with varying  $E_T$  cut
- 3) missing  $E_T$  greater than 20 or 30 GeV


Event selection was optimized for different point in the mSUGRA parameter space.  
However typically  $\epsilon_B \sim 0.05\%$

The background sources


- Z and  $t\bar{t}$  production
- instrumental backgrounds

The numbers of observed events are consistent with those expected from background processes

The null results are interpreted within the mSUGRA framework



95% C.L. Excluded Region



# Charginos and Neutralinos

Production of  $\tilde{\chi}_1^\pm \tilde{\chi}_2^0$  will lead to trilepton events with  $E_T$ , one of the cleanest signature for supersymmetry

Both CDF and DØ searched for trilepton ( $e, \mu$ ) events in Run I



*PRL 80, 1591 (1998); 80, 5275 (1998)*

- 1)  $\geq 3\ell$  ( $e, \mu$ ) with  $E_T > 5-11$  GeV
- 2) require  $e^+e^-$  or  $\mu^+\mu^-$  (CDF only)
- 3)  $E_T > 10-15$  GeV
- 4)  $\ell\ell$  topology and mass cuts

No events were observed in either experiment. CDF expected to see  $1.2 \pm 0.2$  events while DØ estimated to have  $1.3 \pm 0.4$  events from background processes

The null results were interpreted in the framework of MSSM models which give  $m_{\tilde{\chi}_1^\pm} \approx m_{\tilde{\chi}_2^0} \approx 2m_{\tilde{\chi}_1^0}$

The efficiency is typically 3-12% for CDF and 2-6% for DØ when chargino mass is varied from 50 to 100 GeV



# Single-Photon Final State

**DØ (99 pb<sup>-1</sup>) searched for single-photon events with two or more jets and large missing E<sub>T</sub>**

*Phys. Rev. Letters 82, 29 (1999)*

**Within the MSSM, the radiative decay  $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 + \gamma$  dominates in some regions of parameter space**

- 1)  $E_T^\gamma > 20$  GeV,  $|\eta^\gamma| < 1.1$  or  $1.5 < |\eta^\gamma| < 2.5$
- 2)  $\geq 2$  jets with  $E_T^j > 15$  GeV,  $|\eta^j| < 2.0$
- 3)  $H_T$  and  $E_T$  requirements to reduce backgrounds


**Production of supersymmetric particles at Tevatron would lead to anomalous events of photons with large missing E<sub>T</sub>**

**Major background from missing E<sub>T</sub> mismeasurement**

**The missing E<sub>T</sub> distributions of signal and control samples agree very well**


**For mE<sub>T</sub> > 25 GeV, 318 events were selected with 320 ± 20 events expected**

**The null results are interpreted assuming  $p\bar{p} \rightarrow \tilde{q} / \tilde{g} \Rightarrow \tilde{\chi}_2^0 + X \Rightarrow \gamma jj E_T + X$**



# R-parity Violation Signatures


In addition to the SM interactions, following interactions are allowed



resulting lepton and baryon number violations as well as the R-parity violation


**B-violating  $\lambda''_{ijk}$  couplings will lead to multijet events without  $E_T$**

**The L-violating  $\lambda_{ijk}$  and  $\lambda'_{ijk}$  couplings will give rise to multilepton events**



$$\tilde{\chi}_1^0 \rightarrow \nu \tilde{\nu}^* \Rightarrow \nu \ell \ell \quad (\lambda_{ijk})$$

$$\tilde{\chi}_1^0 \rightarrow \ell \tilde{\ell}^* \Rightarrow \ell q q \quad (\lambda'_{ijk})$$



**Frequent assumptions:**

- 1) R-parity violating LSP decay
- 2) couplings are not too weak or too strong
- 3) terms with similar event topology dominate

**Signatures:**

$$p\bar{p} \rightarrow SUSY \Rightarrow \ell^n + j^m (+E_T)$$

# Dilepton ( $\pm\pm$ ) from R-parity Violations

Like-sign dileptons are expected from

$$p\bar{p} \rightarrow \tilde{g}\tilde{g} \rightarrow (\bar{c}\tilde{c}_L)(\bar{c}\tilde{c}_L) \Rightarrow \bar{c}\bar{c}(e^\pm d)(e^\pm d)$$



$$p\bar{p} \rightarrow \tilde{q}\tilde{q} \rightarrow (\bar{q}\tilde{\chi}_1^0)(q\tilde{\chi}_1^0) \Rightarrow q\bar{q}(q\bar{q}'e^\pm)(q\bar{q}'e^\pm)$$

CDF: *Phys. Rev. Lett.* **83**, 2133 (1999)

Motivated in part by HERA excess and probes  $\lambda'_{121}$  R-parity violating coupling

- 1)  $e^\pm e^\pm$  with  $E_T > 15$  GeV
- 2)  $\geq 2$  jets with  $E_T > 15$  GeV
- 3) no significant  $E_T$

No event observed while  $0.3 \pm 0.3$  background events expected



# Dilepton from R-parity Violations

Dzero Studied the case that all RPV couplings are small except  $\lambda'_{1jk}$  within the framework of mSUGRA

DØ: *Phys. Rev. Lett.* **83**, 4476 (1999)



$$p\bar{p} \rightarrow SUSY \Rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow eejjjj$$

- 1)  $E_T^{e1} > 15$  GeV,  $|\eta| < 1.1$  or  $1.5 < |\eta| < 2.5$
- 2)  $E_T^{e2} > 10$  GeV,  $|\eta| < 1.1$  or  $1.5 < |\eta| < 2.5$
- 3)  $\geq 4$  jets with  $E_T^j > 15$  GeV,  $|\eta| < 2.5$
- 4)  $M_{ee}$  not consistent with  $M_Z$

Backgrounds are dominated by Drell-Yan process and mismeasurement


Two events observed with  $1.8 \pm 0.4$  events expected from backgrounds

The analysis assumes  $\tilde{\chi}_1^0$  LSP and only the LSP decays through R-parity violating interactions



# Trilepton from R-parity Violations

The lepton number violating terms in the R-parity violating Lagrangian will result events with four-leptons



DØ has reinterpreted the trilepton analysis of charginos and neutralinos searches in the RPV framework


The analysis assumes

- 1)  $\tilde{\chi}_1^0$  is the LSP
- 2) the LSP decays through RPV

The analysis is sensitive to


- 1) the 1st and 2nd lepton family
- 2) couplings  $10^{-4} < \lambda_{ijk} < 10^{-3}$

The sensitivity is measured using mSUGRA model assuming R-parity violations



# Four-lepton from R-parity Violations

CDF searched four-lepton events from R-parity violating process with a dominant  $\lambda_{121}$  in the mSUGRA framework




$$p\bar{p} \rightarrow SUSY \Rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow llll + X$$

- 1)  $E_T^l > 12, 5, 5, 5$  GeV
- 2)  $\Delta R^{ll} > 0.4$

One event observed while  $1.3 \pm 0.4$  events expected from SM (bb, cc, ...) and instrumental backgrounds

There were 185 tri-lepton events in the data

Efficiency for the signals varies between 10–20% for most of the parameter space of interest



# Gauge Mediation Signatures

Signatures generally depend on the next-lightest supersymmetric particle (NLSP)

$$\begin{aligned}\tilde{\chi}_1^0 &\rightarrow \gamma\tilde{G}, Z\tilde{G}, h\tilde{G} \\ \tilde{l} &\rightarrow lG\end{aligned}$$

$$\begin{aligned}p\bar{p} &\rightarrow SUSY \Rightarrow 2NLSP + \ell^n + j^m \\ &\Rightarrow (E_T + \ell^n + j^m) + \gamma\gamma \\ &\Rightarrow (E_T + \ell^n + j^m) + ll \\ &\Rightarrow (E_T + \ell^n + j^m) + \gamma h \\ &\dots\dots\end{aligned}$$

Signatures:

$$\gamma E_T, ll E_T, \gamma b\bar{b} E_T, \dots$$

Depending on their lifetimes, NLSPs can decay at the production vertex, inside and outside detector



displaced photons  
hot cells  
slow moving particles  
kinked tracks

Only two Run I analyses specifically done for GMSB models:

- light gravitino production
- diphoton+mEt events



# Monojet from Gravitino Production

If the gravitino is light and all other super-partners are heavy, it could be the only super-partner produced at Tevatron, resulting monojet events with large  $m_{E_T}$



- 1) leading jet with  $E_T > 80$  GeV
- 2) at least one central jet with  $E_T > 10$  GeV
- 3)  $E_T > 50$  GeV
- 4) no high  $p_T$  leptons
- 5) topological cuts to reduce mismeasurement backgrounds

**Backgrounds are dominated by W/Z+jets production. No excess was observed.**



**19 events were observed with  $m_{E_T} > 175$  GeV while 22 were expected. This leads to**

$$\sqrt{F} > 217 \text{ GeV} \Rightarrow m_{\tilde{G}} \geq 1.1 \times 10^{-5} \text{ eV}$$


# Diphoton from Neutralino Decays

Both CDF and Dzero searched for  $p\bar{p} \rightarrow \gamma\cancel{E}_T + X$  events and no significant excess was found


CDF: PRD 59, 092002 (1999) DØ: PRL 80, 442 (1998)

They are expected from

$p\bar{p} \rightarrow SUSY \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0 + X$  with  
 $\tilde{\chi}_1^0 \rightarrow \gamma + \tilde{G}$  decays




Events with two photons and large missing  $E_T$  are rare



# Large Extra Dimensions

The existence of large extra dimensions will modify both the mass and the angular distributions of dilepton and diphoton events




DØ has searched for virtual graviton effects in the final state with two electromagnetic clusters, combining dielectron and diphoton final states in the analysis to eliminate cross Misidentifications between electrons and photons

$$p\bar{p} \rightarrow (ee, \gamma\gamma) + X$$

$$\frac{d^2\sigma}{dM d\cos\theta^*} = f_{SM}(M, \cos\theta^*) + f_{\text{int}}(M, \cos\theta^*)\eta + f_{KK}(M, \cos\theta^*)\eta^2$$

where  $\eta = \frac{F}{M_S^4}$

MC Simulation of the ED signatures



# Dielectron+Diphoton Final States


- 1) two electromagnetic clusters with  $E_T > 45$  GeV,  $|\eta| < 1.1$  or  $1.5 < |\eta| < 2.5$
- 2)  $E_T < 25$  GeV

The non-separation between electrons and photons yield very high efficiency (76%) for di-EM final states

1282 events were selected from a sample of  $127 \text{ pb}^{-1}$

Comparison of the data and the SM predictions


DØ Preliminary, Run I,  $127 \text{ pb}^{-1}$



Background sources:

- Drell-Yan dielectron
- $\gamma\gamma$  production
- instrumental

Comparison of the data with the SM predictions



# Limits on LED Scale

The kinematic distributions of di-EM final states are well described by the sum of the SM and instrumental backgrounds

**No evidence for large extra dimensions**

The leading-order distributions were augmented with


- initial state radiation effect with a transverse kick
- K-factor correction to account for the NLO effect.

The total systematic error is estimated to be about 19%.

Binned maximum likelihood fit to extract the limits.

$M_s > 1.3$  TeV ( $n = 2$ )  
1.4 TeV ( $n = 3$ )  
1.2 TeV ( $n = 4$ ) @ 95% C.L.

Limits on Large Spatial Extra Dimensions



# Run II Upgrade & Schedule

## Major Tevatron Improvements


- 1) Replace main ring with main injector
- 2) Construct a new anti-proton storage ring
- 3) Collider center-of-mass energy of  $\sim 2$  TeV

The number of anti-protons in the ring has been one of the major limiting factors in Tevatron luminosity. The anti-proton stacking rate will be increased to  $2 \times 10^{11}/\text{hr}$  from  $7 \times 10^{10}/\text{hr}$

The machine will operate with 36x36 bunches (396 ns spacing) Initially and with 140x121 bunches (132 ns) eventually.



## Run II machine goals:

- 1) Run IIa to achieve a luminosity of  $5 \times 10^{31} \text{ cm}^{-2}\text{s}^{-1}$  and an integrated luminosity of  $2 \text{ fb}^{-1}$
- 2) Run IIb to achieve a luminosity of  $2 \times 10^{32} \text{ cm}^{-2}\text{s}^{-1}$  and an integrated luminosity of  $\sim 20 \text{ fb}^{-1}$



**Run II comes to us  
March 1, 2001**

# CDF and DØ Detectors (Run II)



- a new massive silicon vertex detector
  - 7 layers extending to 28 cm in radius
  - dead timeless SVX3 readout electronics
- a new central outer tracker
- fast, hermetic scintillator tile plug and forward calorimeter
- large trigger bandwidth

- entirely new tracking system
  - 2T super conductor solenoid
  - disk/barrel silicon detector
  - 8 layers of scintillating fiber tracker
  - preshower detectors
- improved muon spectrometer
- new trigger and DAQ system

# Squark and Gluino in Run II

**Squarks and gluinos will remain to be copiously produced if kinematically accessible**

- **Though leptonic events are also expected, the final state with multijet and large mEt remains to be the dominant signature of squark and gluino production except for isolated regions of SUGRA parameter space.**
- **For large  $\tan\beta$  values, gluino, chargino and neutralino decays to third generation particles are significantly enhanced.**
- **Better understanding of mEt tail of multijet events and developing a robust method to accurately estimate multijet backgrounds are critical for squark and gluino searches in Run II.**
- **It is important to have good tau-lepton and b-quark trigger and identification capabilities.**

## **Run II improvements:**

- 1) **mEt resolution improvement**
  - more hermetic detector (CDF)
  - better vertexing (DØ)
- 2) **Advanced analysis methods and generally improved tools**

**With  $2\text{fb}^{-1}$ , the upgraded Tevatron should be able to probe  $m_{1/2}$  up to  $\sim 150$  GeV, corresponding to a gluino mass of  $\sim 400$  GeV if  $m_0 < 200$  GeV**

# Chargino and Neutralino in Run II

In mSUGRA model,  $\tilde{\ell}, \tilde{\chi}_1^\pm, \tilde{\chi}_1^0, \tilde{\chi}_2^0$  are typically less massive than  $\tilde{q}, \tilde{g}$ .

So the  $p\bar{p} \rightarrow \tilde{\chi}_1^\pm \tilde{\chi}_2^0 + X$  production is one of the most promising channels for the searches at Tevatron

At large  $\tan\beta$ , the lighter tau slepton  $\tilde{\tau}_1$  is lighter than  $\tilde{\chi}_2^0, \tilde{\chi}_1^\pm$ . Then,  $\tilde{\chi}_2^0$  and  $\tilde{\chi}_1^\pm$  can dominantly decay into final states with  $\tau$  via  $\tilde{\tau}_1$

The signature is therefore trilepton events with large tau content


Major backgrounds:

W+jets, Z+jets, WZ,...

(WZ background is irreducible)

Key to success:

large acceptances and high efficiencies for high and low  $p_T$  leptons including the tau lepton






Run II improvements:

- 1) extended detector coverage for leptons to improve acceptances
- 2) lepton charge measurement and better muon momentum measurement to reduce backgrounds (DØ)
- 3) major effort on tau identification built upon Run I experiences


For small to medium values of  $\tan\beta$  values, maximum chargino mass reach of  $\sim 200 \text{ GeV}$  is achievable at Tevatron Run II

# Long-lived Neutralino


The upgraded DØ detector will enable us to identify non-vertex pointing photons by matching clusters in the EM calorimeter and preshower detectors



Providing unique tools in searching for long-lived neutralinos





# Long-lived Slepton



## Tools for massive stable charged particles (MSP)

dE/dx information from

- 1) Silicon
- 2) Fiber tracker
- 3) Preshowers
- 4) Calorimeter



An efficiency of 68% for MSP and a rejection factor of 10 for MIP are assumed

# Summary

**We have carried out extensive searches for physics beyond standard model in Run I. Nothing against or for supersymmetry has been found.**



**We will continue the searches in Run II and expect to explore gluino mass up to 400 GeV and chargino mass up to 200 GeV.**



**To make discovery, we need help, not only from theorists, but also from the God.**

